



NOTICE

Before using this product, read the entire label

Bortrac[®] 150

YaraVita[™]

Bortrac[®] 150

EC FERTILISER

Inorganic foliar fertiliser

Guaranteed analysis:

15% w/v = **150g/l boron (B)** [10.9% w/w] soluble in water as boron ethanalamine

pH:8

To be used only where there is recognised need.
Do not exceed the appropriate application rates.

Batch number: see pack for details.

Manufacture date: See pack for details.

Validity date: 5 years from date of manufacture.

H402: Harmful to aquatic life.

P273: Avoid release to the environment.

5 litres

Net Weight
6.76Kg

Distributed by: **Yara East Africa Ltd.**

P.O. Box 50949, City Square 00200, Nairobi, Kenya.

Tel: +254 0724 255 370 www.yara.co.ke

YaraVita™ Bortrac® 150

APPLICATION RATES AND TIMINGS

Aubergine, Cucumber (Field Grown): 1 to 2 l/ha as soon as there is sufficient foliage to intercept spray. A second application may be made at the same rate 10 to 14 days later. Water rate: 50 to 200 l/ha.

Avocado: 2 l/ha at spring bud development and again at spring flush. Water rate: minimum 500 l/ha.

Bananas: 1 l/ha. Regular applications may be necessary where low or marginal levels of boron exist. Water rate: minimum 30 l/ha.

Brassicas, Broccoli, Cabbage, Chinese Cabbage: 3 l/ha at 4 to 6 leaf stage with repeat applications at the above rate at 10 to 14 day intervals for moderate to severe deficiency. Water rate: minimum 50 l/ha.

Cereals (Barley, Rice, Wheat): 1 l/ha applied at the start of tillering. Water rate: 30 to 200 l/ha.

Coffee: 1 l/ha early in the growing season, pre-flowering and again post-harvest. Water rate: minimum 200 l/ha.

Groundnuts/Peanuts: 2 l/ha at the 4 to 6 leaf stage. Water rate: minimum 50 l/ha.

Maize, Sorghum: 3 l/ha at 4 to 8 leaf stage. For moderate to severe deficiency, a repeat application may be necessary 10 to 14 days later. Water rate: minimum 30 l/ha.

Peas (Field Grown): 2 l/ha at 4 to 6 leaf stage. For moderate to severe deficiency, a repeat application may be necessary 10 to 14 days later. Water rate: minimum 200 l/ha.

Pepper (Field Grown): 2 l/ha applied at early flowering to fruiting, with two repeat applications at 10 to 14 day intervals if necessary. Water rate: minimum 200 l/ha.

Pineapple: 1 to 2 l/ha pre-flowering. Water rate: minimum 200 l/ha.

Squash (Field Grown): 2 l/ha from the 4 leaf stage. Repeat at 10 to 14 days intervals if necessary. Water rate: minimum 50 l/ha.

Sugar Cane: 2 l/ha when cane is between 30 and 120 cm tall. Repeat applications may be necessary at 10 to 14 day intervals. Water rate: minimum 30 l/ha.

Tea: 0.5 l/ha applied during plucking season. Repeat at approximately 1 month intervals. Spray 2 to 6 applications. Water rate: minimum 200 l/ha.

Tobacco: Two applications of 2 l/ha two to three weeks after transplanting (3 to 4 leaf stage) with 10 days between applications. Water rate: minimum 30 l/ha.

Vegetables [Beans, Garlic, Leek*, Onions, Potatoes, Tomatoes (Field Grown)]: 1 to 2 l/ha as soon as there is sufficient foliage to intercept a spray. Repeat 10 to 14 days later if necessary. Water rate: 200 l/ha. *Do not apply for two weeks after transplanting.

FERTIGATION: Use at 0.3 litres per 1000 metres squared. Repeat at 7 to 14 day intervals as necessary.

PROTECTED CROPS: 0.1 litres per 100 litres water maximum concentration. Water rate: 1000 l/ha maximum. Refer to equivalent field grown crop for application timing.

For further crop recommendations contact your local Yara representative.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the SDS for further information, available at www.yara.com.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.