



**YaraVita**<sup>™</sup>  
**Caltrac**<sup>™</sup>



#### NOTICE

Before using this product, read the entire label

# Caltrac<sup>™</sup>

## EC FERTILISER

Mixture of micro-nutrients boron (B) and zinc (Zn) with calcium (Ca) and nitrogen (N)

Guaranteed analysis:

4.1% w/w = 69 g/l Nitrogen (N) as urea

23.2% w/w = 400 g/l Calcium (Ca),

0.2% w/w = 3 g/l Boron (B),

1.8% w/w = 31 g/l Zinc (Zn),

Maximum biuret: 0.1%w/w

To be used only where there is a recognised need.  
Do not exceed the appropriate application rates.

Manufactured by: Yara UK Ltd., Pocklington, York,  
YO42 1DN, UK  
Tel: 01759 302545  
Distributed by: **Yara International (M) Sdn Bhd**,  
Lot 3.02, Level 3, 1 First Avenue, Bandar Utama  
47800 Petaling Jaya, Selangor Darul Ehsan,  
Malaysia  
Tel: 603 76138010

1 litre

Net weight  
1.72 kg

#### RECOMMENDED APPLICATION RATES AND TIMINGS

**Apple, Pears:** 3-5 L/ha repeated at 10 day intervals from 4 weeks after petal fall. Water rate: 500 L/ha minimum.\*

**Avocado:** Apply at a concentration of 150 mLs product per 100 litres water from fruit set at 10 to 14 day intervals to a maximum tree row volume (TRV) of 4000 L/ha.\*

**Bananas:** 2 L/ha repeated as necessary at 10 to 14 day intervals. Water rate: 30 L/ha minimum.\*

**Brassicac, Canola:** 3 to 5 L/ha applied from the 4 to 9 leaf stage. Repeat as necessary at 10 to 14 day intervals. Water rate: 50-500 L/ha.

**Capsicums (Field Grown):** Repeat applications of 5 L/ha from the 4 to 6 leaf. Allow 7 days between applications.

Water rate: 50-500 L/ha.\*

**Cucurbits (Field Grown):** 3 to 5 L/ha repeated at 7 day intervals during fruit development. Water rate: 200-500 L/ha.\*

**Grapevines:** Two applications of 5 L/ha at a 10 to 14 day interval prior to flowering. Also, repeated applications of 5 L/ha at 10 to 14 day intervals from fruit set e.g. fruit set, bunch closure (pea sized berries), start of ripening and two weeks later. Water rate: 200 L/ha. NB Late applications may leave a deposit on berries which may be of significance for table grapes.

**Legumes/Pulses:** 3 to 5 L/ha at the 4 to 6 leaf stage. For moderate to severe deficiency repeat at 10 to 14 day intervals, as necessary. N.B. Do not apply during flowering. Water rate: 30 to 200 L/ha.

**Onion:** 3 to 5 L/ha applied at the 6 leaf stage and repeated as necessary 10 to 14 days later. Also, 5 L/ha at bulb swelling, repeated once or twice at 10 to 14 day intervals. Water rate: 30-200 L/ha.

**Potatoes:** 5 L/ha commencing 2 weeks after 100% emergence and repeated as necessary at 10 to 14 day intervals. Water rate: 50-500 L/ha.

**Stonefruit:** Two applications of 5 L/ha sprayed 6 weeks and 4 weeks before harvest. Water rate: 500 L/ha. Alternatively: Regular applications of 5 L/ha at 7 to 14 day intervals from petal fall to one month before harvest. Note: Final application to be made at least one month before harvest. Under conditions conducive to fruit russet (variety, weather conditions, etc) delay applications until 6 weeks after petal fall. Water rate: 150 to 500 L/ha.

**Tomato (Field Grown):** One to four applications of 5 L/ha from flowering on second truss onwards, repeated at 7 day intervals. Water rate: 50-500 L/ha.\*

**Zucchini (Field Grown):** 3 to 5 L/ha repeated at 7 day intervals during fruit development up to one month before harvest. Water rate: 200-500 L/ha.\*

\*Note: Caltrac 400 may leave white marks upon the fruit, similar to other suspension concentrate products. It may be necessary to clean the fruit prior to sale.

#### DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS  
Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

#### TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.

#### PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

#### STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 40°C) Dispose of contents and container in accordance with all local, regional, national and international regulations.

#### CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

H401: Toxic to aquatic life.

H412: Harmful to aquatic life with long lasting effects.

P273: Avoid release to the environment.