



NOTICE

Before using this product, read the entire label

Caltrac BZ™

YaraVita™

Caltrac BZ™

GROUP 2 FERTILIZER

REGISTRATION No. B3715 Act 36 of 1947

Mixture of trace elements boron and zinc with calcium

To be used only where these is recognised need.
Do not exceed the appropriate application rates.

Guaranteed analysis:

2g/kg = 3g/l boron (B) total,

18g/kg = 31g/l zinc (Zn) total,

232g/kg = 400g/l calcium (Ca) total,

Typical density 1.72kg/litre at 20°C

H401: Toxic to aquatic life.

H412: Harmful to aquatic life with long lasting effects.

P273: Avoid release to the environment.

10 litres

Net Weight
17.24kg

Manufactured by Yara UK Ltd., Pocklington, York YO42 1DN, U.K.

Registered by Yara Africa Fertilizers (PTY) Ltd.,

Reg.nr. 2011/003434/07

The Pivot @ Montecasino, No. 1 Montecasino Boulevard, Fourways,
Johannesburg 2191.

YaraVita™ Caltrac BZ™

RECOMMENDED APPLICATION RATES AND TIMINGS

Apples: 3 to 5 l/ha repeated at 7 to 10 day intervals applied from petal fall until 6 weeks before harvest. Water rate: 500 l/ha minimum.

Avocado: Up to 300 mls/100 l from fruit set until one month before harvest. Based on a maximum TRV of 2,000 l/ha.

Citrus: 5 l/ha from fruit set until 6 weeks before harvest at 10 to 14 day intervals. Water rate: 500 l/ha minimum. Alternatively, 150 mls / 100 l from fruit set until 6 weeks before harvest at 10 to 14 day intervals, based on a typical TRV of 2500 l/ha and a maximum TRV of 4000 l/ha.

Field Vegetables (Broccoli, Brussel Sprouts, Cabbage, Calabrese, Cauliflower, Carrots, Lettuce): 3 to 5 l/ha from the 4 to 9 leaf stage.

Repeat as necessary at 10 to 14 day intervals until one month before harvest. Water rate: 50 to 500 l/ha.

Grapevines: Two applications of 5 l/ha at a 10 to 14 day interval starting from fruit set. Water rate: 200 l/ha. N.B: Late applications may leave a deposit on berries which may be of significance for table grapes.

Groundnuts/Peanuts: 3-5 l/ha applied at 4-6 leaves.

Water rate: 30-200 l/ha

Mango: 5 l/ha. Apply one to two sprays 7 to 10 days apart during new leaf flush. Do not apply during flowering. 5 l/ha from fruit set at 10 to 14 day intervals. N.B. Late applications may leave a deposit on the fruit. Water rate: 500 l/ha minimum.

Melon, Cucumber, Squash, Pumpkin: Repeat applications of 2 l/ha at 7-10 day intervals from the first fruit set up to one month before harvest. Water rate: 200 to 400 l/ha.

Nuts (Deciduous): 5 l/ha 7 days after petal fall. Water rate: 500 l/ha.

Pears: 3 to 5 l/ha repeated at 7 to 10 day intervals from petal fall until 6 weeks before harvest. Water rate: 500 l/ha minimum. Alternatively, 150 mls / 100 litres repeated at 7 to 10 day intervals from petal fall until 6 weeks before harvest. Based on a typical TRV of 2500 l/ha and a maximum TRV of 4000 l/ha.

Pepper (Field Growth): Repeat applications of 5 l/ha from the 4 to 6 leaf stage up to one month before harvest. Allow 7 days between applications. Water rate: 50 to 500 l/ha.

Stone Fruit (Apricot, Cherry, Nectarines, Peaches, Plums): Regular applications of 5 l/ha at 7 to 14 day intervals from petal fall until 6 weeks before harvest. Water rate: 500 l/ha minimum. Alternatively, regular applications of 125 mls/100 litres water at 7 to 14 day intervals from petal fall to 6 weeks before harvest. Based on a typical TRV of 2,500 l/ha and a maximum TRV of 4,000 l/ha. Note: Under conditions conducive to fruit russet (variety, weather conditions, etc) delay applications until 6 weeks after petal fall.

Strawberry (Field Growth): 3 to 5 l/ha applied three times from start of new season leaf growth at 10 to 14 day intervals. N.B. Last application to be made no later than one month before harvest commences. Also, 3 to 5 l/ha applied as required after final harvest for the season at 10 to 14 day intervals. Water rate: 200 to 500 l/ha.

Tomato (Field Growth): One to four applications of 5 l/ha from flowering on second truss onwards up to one month before harvest. Allow 7 days between applications. Water rate: 50 to 500 l/ha.

For further crop recommendations contact your Yara representative.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the MSDS for further information, available at www.yara.com.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

