



YaraVita™

Brassitrel™



NOTICE

Before using this product, read the entire label

Brassitrel™

EC FERTILISER

Mixture of micro-nutrients boron (B), manganese (Mn) and molybdenum (Mo) with nitrogen (N), magnesium (Mg) and calcium (Ca)

Guaranteed analysis - total:

4.5% w/w = **69 g/l Nitrogen (N)**

[4.5% ureic nitrogen]

4.6% w/w = **70 g/l Magnesium (Mg)**

5.8% w/w = **89 g/l Calcium (Ca)**

3.9% w/w = **60 g/l Boron (B)**

4.6% w/w = **70 g/l Manganese (Mn)**

0.3% w/w = **4 g/l Molybdenum (Mo)**

To be used only where there is a recognised need.
Do not exceed the appropriate application rates.

Manufactured by: Yara UK Ltd., Pocklington, York, YO42 1DN, UK

Tel: 01759 302545

Distributed by: **Yara International (M) Sdn Bhd**, Lot 3.02, Level 3, 1 First Avenue, Bandar Utama 47800 Petaling Jaya, Selangor Darul Ehsan, Malaysia

Tel: 603 76138010



Warning

Hazardous ingredients: colemanite (calcium borate), 2-methylisothiazol-3(2H)-one

H317: May cause an allergic skin reaction. H361: Suspected of damaging fertility or the unborn child. P202: Do not handle until all safety precautions have been read and understood. P280: Wear protective gloves/ clothing and eye/ face protection. P261: Avoid breathing gas, vapor or spray. P302+P352: IF ON SKIN: Wash with plenty of soap and water. P308+P313-a: IF exposed or concerned: Get medical attention.

5 litres

**Net weight
7.68kg**

YaraVita Brassirel™

APPLICATION RATES AND TIMINGS

Beans, Peas: Up to 4 l/ha as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4 to 6 leaf stage). Repeat as necessary at 10 to 14 day intervals up to one month before harvest. N.B. Do not spray during flowering. Water rate: 200 l/ha.

Broccoli, Brussel Sprouts, Cabbage, Calabrese, Cauliflower: Up to 4 l/ha as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4 to 6 leaf stage). Repeat as necessary at 10 to 14 day intervals up to one month before harvest. Water rate: 200 l/ha.

Carrot: Two applications of 3 l/ha applied when the crop is 15 cm tall and again 2 weeks later. Water rate: 200 l/ha.

Cereals: 3 l/ha applied at any time from the 2 leaf stage to 1st node detectable (Zadoks G.S. 12 to 31). Water rate: 200 l/ha.

Leek: 3 l/ha applied two weeks after transplanting or in the case of direct sown crops, when the crop is 15 cm tall. For moderate to severe deficiency one or two repeat applications may be necessary at 10 to 14 day intervals. Water rate: 200 l/ha.

Linseed: 3 l/ha when crop is 15 cm tall. For moderate to severe deficiency repeat applications at 10 to 14 day intervals. Water rate: 200 l/ha.

Lupin: 3 l/ha at the 4 to 6 leaf stage. Repeat once or twice at 10 to 14 day intervals if necessary. (For winter varieties apply the repeat sprays at spring re-growth). Water rate: 200 l/ha.

Oilseed Rape: 3 to 4 l/ha at the 4 to 9 leaf stage and again at the onset of the rapid stem extension phase up to the start of flowering. Repeat at 10 to 14 day intervals within these growth stages as necessary. Also, 3 l/ha at the end of petal fall. Water rate: 50 to 200 l/ha.

Onion: 2 applications of 3 l/ha applied as soon as there is adequate leaf cover (6 leaves) and repeated 2 weeks later. Water rate: 200 l/ha.

Potatoes: 3 l/ha applied one week after 100 % emergence with a second application 10 to 14 days later. Water rate: 200 l/ha.

Soya Bean: Up to 4 l/ha as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4 to 6 leaf stage). Repeat as necessary at 10 to 14 day intervals up to one month before harvest. N.B. Do not spray during flowering. Water rate: 200 l/ha.

Sugar Beet: 3 l/ha in 200 litres of water. Repeat applications can be made at 10 to 14 day intervals. Apply from 4 to 6 leaf stage.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS
Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 40°C)
Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.