



**YaraVita™**

**Coptrac™ 500**



**5 litres**

**Net Weight  
7.62kg**

## NOTICE

Before using this product, read the entire label

# Coptrac™ 500

GROUP 2 FERTILIZER

REGISTRATION No. B3713 Act 36 of 1947

### Copper oxide suspension

To be used only where these is recognised need.  
Do not exceed the appropriate application rates.

Guaranteed analysis:

**330g/kg = 500g/l copper (Cu) total**

Typical density 1.52kg/litre at 20°C

**Contains compounds of copper. Do not breath spray.**

Manufactured by Yara UK Ltd., Pocklington, York YO42 1DN, U.K.  
Registered by Yara Africa Fertilizers (PTY) Ltd.,  
Reg.nr. 2011/003434/07  
The Pivot @ Montecasino, No. 1 Montecasino Boulevard,  
Fourways, Johannesburg 2191.

## Danger

Contains: dicopper oxide  
H302: Harmful if swallowed.  
H318: Causes serious eye damage. H410:Very toxic to aquatic life with long lasting effects.P280: Wear protective clothing and eye protection.  
P270: Do not eat, drink or smoke when using this product.  
P305+P351+P338+P310: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.  
Immediately call a POISON CENTER or doctor/physician.  
P301+P312: IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell. P391: Collect spillage.



**UN 3082**  
ENVIRONMENTALLY  
HAZARDOUS SUBSTANCE,  
LIQUID, N.O.S. (dicopper oxide)

# YaraVita™ Coptrac™ 500

## RECOMMENDED APPLICATION RATES AND TIMINGS

**Apples, Pears:** Based on a typical TRV (tree row volume) of 2500 l/ha, apply 20 mls / 100 litres of water, to a maximum of 0.5 l/ha post-harvest, before leaf fall.

**Avocado:** 0.5 l/ha applied to flush of new growth prior to flowering. Water rate: 500 to 1000 l/ha.

**Canola:** 0.25 l/ha at the 4 to 8 leaf stage. Water rate: 200 l/ha.

**Cereals:** 0.25 to 0.5 l/ha at tillering to second node detectable (Craven 2 - 15). Water rate: 50 to 200 l/ha. For moderate to severe deficiency, repeat applications at 10 to 14 day intervals.

For aerial application apply 0.25 l/ha at the timings described above. Water rate: 20 to 50 l/ha.

**Seed Dressing:** Typically 3 litres per tonne. Also, up to 6 litres per tonne if necessary.

**Citrus:** 200 mls / 100 litres water to give a maximum of 2 l/ha / application post harvest. Not to be used when either flowers or fruit are on the tree. Alternatively, 0.5 l/ha applied to both spring and autumn flushes. Water rate: 500 l/ha minimum.

**Grapevines:** 0.5 l/ha before flower buds visible stage.

Water rate: 200 to 1000 l/ha.

**Maize:** 0.5 l/ha at the 4 to 8 leaf stage. Water rate: 75 to 200 l/ha.

Seed Dressing: 4 litres per tonne.

**Olive:** Spray at 1.25 to 2.5 l/ha. Two to three sprays per year between spring and end of autumn. To be applied when problem is apparent. 10 to 14 days between applications. Water rate: 1000 l/ha.

**Onion:** 0.5 l/ha two weeks after transplanting, or in the case of direct sown crops, when the crop is 15 cm tall. Repeat applications at 10 to 14 day intervals. Water rate: 75 to 200 l/ha.

**Potatoes:** 0.5 l/ha applied 7 to 14 days after 100% emergence and following petiole analysis during tuber bulking.

Water rate: 200 l/ha (30 l/ha for aerial application).

**Stone Fruit (Apricot, Cherry, Nectarine, Peach, Plum):** 0.5 l/ha applied after harvest before leaf fall. Water rate: 1000 l/ha.

Alternatively for South Africa: Based on a typical TRV (tree row volume) of 2500 l/ha, apply 20 mls / 100 litres of water, to a maximum of 0.5 l/ha post-harvest, before leaf fall.

**Sugar Cane:** 0.5 l/ha when cane is between 30 and 120 cm tall. Repeat applications may be necessary at 10 to 14 day intervals. Water rate: 30 to 200 l/ha.

**Tomato (Field grown):** 0.5 l/ha at the 4 to 6 leaf stage, (or 35 days after transplanting). Water rate: 200 l/ha.

**For further crop recommendations contact your Yara representative.**

## DIRECTIONS FOR USE

### ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

### TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.

## PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the MSDS for further information, available at [www.yara.com](http://www.yara.com).

## STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

## CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

