

YaraVita[¬]

Crop Boost[™]



NOTICE

Before using this product, read the entire label

Crop Boost[™]

EC FERTILISER

Inorganic foliar fertiliser PK (Mg) fertiliser solution 29.5 - 5 (2.7) with zinc (Zn)

Guaranteed analysis - Soluble in water:

29.5% = 440g/l phosphorus pentoxide (P,O,) [12.9% phosphorus (P)], 5% = 75g/l potassium oxide (K,O) [4.2% potassium (K)],

4.5% magnesium oxide (MgO) [2.7% = 40g/l Magnesium (Mg)],

3.1% = 46g/l zinc (Zn)

pH:1

Manufacture date: See top of pack. Expiry date: 5 years from date of manufacture.

Distributed by: Yara East Africa Ltd. P.O. Box 50949, City Square 00200, Nairobi, Kenya. Tel: +254 0724 255 370 www.vara.co.ke Distributed by: Yara Tanzania Ltd. P.O.Box 40230 Dar es Salaam, Tanzania. Tel: +255 737 22 5802/3 Manufactured in the UK by Yara UK Ltd., Pocklington, York YO42 1DN, U.K. Tel: 00 44 1759 302545

DANGER

Hazardous ingredients: zinc bis(dihydrogen phosphate), phosphoric acid H290: May be corrosive to metals. H303: May be harmful if swallowed. H314: Causes severe skin burns and eye damage. H401: Toxic to aquatic life. H412: Harmful to aquatic life with long lasting effects. P260: Do not breathe gas or vapour. P280: Wear protective gloves/clothing and eye/face protection. P273: Avoid release to the environment. P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P303+P361+P353: IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water. P234: Keep only in original packaging.

CORROSIVE LIQUID, ACIDIC, INORGANIC, N.O.S. (Phosphoric acid)

10 litres

NLTZYP32M00 V.11.24

YaraVita[™] Crop Boost[™]

APPLICATION BATES AND TIMINGS

Cereals (Wheat and Barley): 1 l/ac from the 2 leaf stage to 1st node detectable. Water rate: 80 to 120 l/ac.

Maize: 1 - 2 l/ac at the 4 to 6 leaf stage. Repeat 10 to 14 days later if necessary. Water rate: 80 to 120 l/ac.

Rice: 1 l/ac applied at tillering and again at panicle initiation. Repeat 7 to 10 days later and again after flowering if necessary. Water rate: 80 to 120 l/ac.

Onion: 1-2 l/ac. Apply when the foliage is 15cm tall, with a second application 10 to 14 days later if necessary. Also, 1 to 2 applications of 1l/ac during bulb filling, with a 10 to 14 day interval between sprays. Water rate: 80 to 120l/ac.

Potatoes: To stimulate early growth: 1-2 l/ac one week after 100 percent crop emergence. To increase tuber number: 1-2 l/ac at tuber initiation. To increase tuber size: apply minimum 2 times of 1-2 l/ac during tuber bulking . Allow 10 to 14 days between applications. Water rate: 80-120 l/ac.

Sugarcane: 1-2 l/ac when cane is between 30 and 120 cm tall. Repeat applications may be necessary at 10 to 14 day intervals. Water rate: 80-120 l/ac.

Tomato, Eggplant, Chili, Capsicum (Field Grown): 1-2l/ac applied at the 4 to 6 leaf stage. Repeat at 10 to 14 day intervals as necessary. Maximum 3 applications. Do not apply during flowering. Water rate: 80-120 l/ac.

For further crop recommendations contact your Yara representative.

Kitunguu: Nyunyizia lita 1-2 kwa eka wakati majani yana urefu wa 15cm, rudia baada ya siku 10-14 kama kuna hitaji. Pia nyunyizia mara mbili lita 1 kwa eka wakati wa kunenepa kitunguu kwa mpishano wa siku 10-14. Kiasi cha maji: Lita 80-120 kwa eka.

Mahindi: Nyunyizia lita 1 - 2 kwa eka mahindi yakiwa na majani 4-6. Unaweza kurudia siku 10-14 kama kuna hitaji. Kiasi cha maji: Lita 80-- 120 kwa eka.

Miwa: Nyunyizia lita 1-2kwa eka wakati mua una urefu wa 30-120cm. Rudia kama kuna hitaji kila baada ya siku 10-14. Kiasi cha maji: Lita180-120 kwa eka.

Mpunga: Nyunyizia lita 1 kwa eka mpunga unapoanza kupacha, na wakati wa kutunga mimba dume. Rudia baada ya siku 7 ---10, na baada ya kuchanua kama kuna hitaji. Kiasi cha maji: Lita 80-- 120 kwa eka.

Nafala (Shairi na Ngano): Nyunyizia lita 1 kwa eka mmea ukiwa na majani 2 hadi fundo la kwanza. Kiasi cha maji: Lita 80-- 120 kwa eka.

Nyanya, Biringanya, Pilipili, Hoho (Ukuaji wa Shamba): Nyunyizia lita 1-2 kwa eka wakati mazao yana majani 4-6. Rudia baada ya siku 10-14 kama kuna hitaji. Usinyunyizie zaidi ya mara 3. Usinyunyizie wakati wa maua. Kiasi cha maji: Lita 80-l20 kwa eka.

Viazi Ulava: Kuchochea uotaji wa mapema, nyunyizia lita 1-2 kwa eka wiki moja baada ya mmea kuota kwa asilimia 100. Kwa kuongeza idadi ya viazi, nyunyizia lita 1-2 kwa eka wakati wa viazi vinapoanza kutungwa. Kwa kuongeza ukubwa wa viazi; nyunyizia wastani mara 2 lita 1-2 kwa eka wakati wa kunyanyulia udongo. Zingatia siku 10-14 kabla ya kurudia kunyunyiza. Kiasi cha maji : Lita 80- 120 kwa eka.

DIRECTIONS FOR USE ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraving is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the SDS for further information, available at www.vara.com.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes $(-5^{\circ}C < and < 35^{\circ}C)$. Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

