



**YaraVita™**

**Ferleaf™ 100**



**5 litres**  
**Net weight**  
**6.6 kg**

#### **NOTICE**

Before using this product, read the entire label

# **Ferleaf™ 100**

#### **UK FERTILISER**

Iron fertiliser solution (EDTA)

Guaranteed analysis - soluble in water:  
**7.6% (= 100 g/l) Iron (Fe)** chelated by EDTA

Chelated fraction stable within pH range 1.5 - 6.5

To be used only where there is a recognised need.  
Do not exceed the application rate.

#### **INFORMATION FOR ROI MARKET**

**PFC 1(C)(II)(a) Straight Inorganic micronutrient fertilizer**  
Liquid Solution

Content % w/w:  
**7.6% (= 100 g/l) Iron (Fe)** chelated by EDTA, water soluble

Chelated fraction stable within pH range 1.5 - 6.5

**Ingredients:** Ethylenediaminetetraacetic acid, ferric ammonium complex  
CAS n° 21265-50-9 (CMC 1: Virgin material substances and mixtures).

To be used only where there is a recognised need.  
Do not exceed the application rate.

Batch Number/ Manufacture Date: See Top of Pack

Produced and Distributed by: **Yara UK Limited.**  
Pocklington Industrial Estate, Pocklington, York. YO42 1DN  
Tel: 01759 302545 [www.yara.co.uk](http://www.yara.co.uk)

**ROI:** Imported by: **Yara UK Limited.** The Old Rectory, Cork Road,  
Carrigaline, P43 W921. County Cork.  
Tel: +44 1759 302 545 [www.yara.ie](http://www.yara.ie)

# YaraVita™ Ferleaf™ 100

## APPLICATION RATES AND TIMINGS

**Apple:** 1 litre/ha in 1,000 litres of water, repeated 3-8 times at 7-14 day intervals. Applications should commence as soon as there is sufficient leaf cover (preventative treatment) or as soon as first symptoms are seen (curative treatment).

Do not apply to fruit trees during flowering.

**Cereals:** 1 l/ha in 1,000 litres of water, repeated 3-8 times at 7-14 day intervals. Applications should commence as soon as there is sufficient leaf cover (preventative treatment) or as soon as first symptoms are seen (curative treatment).

Do not apply during flowering.

**Grapevines:** 1 litre/ha in 1,000 litres of water, repeated 3-8 times at 7-14 day intervals. Applications should commence as soon as there is sufficient leaf cover (preventative treatment) or as soon as first symptoms are seen (curative treatment).

Do not apply during flowering.

**Grass (Amenity):** 1 l/ha as soon as growth commences in spring and/or following identification of need by analysis. Repeat sprays at 10 to 14 day intervals as necessary.

Water rate: 200 l/ha.

**Maize:** 1 litre/ha in 1,000 litres of water, repeated 3-8 times at 7-14 day intervals. Applications should commence as soon as there is sufficient leaf cover (preventative treatment) or as soon as first symptoms are seen (curative treatment).

Do not apply during flowering.

**Peas:** 1 litre/ha in 1,000 litres of water, repeated 3-8 times at 7-14 day intervals. Applications should commence as soon as there is sufficient leaf cover (preventative treatment) or as soon as first symptoms are seen (curative treatment).

Do not apply during flowering.

**Plum:** 1 litre/ha in 1,000 litres of water, repeated 3-8 times at 7-14 day intervals. Applications should commence as soon as there is sufficient leaf cover (preventative treatment) or as soon as first symptoms are seen (curative treatment).

Do not apply to fruit trees during flowering.

**Strawberry (Field Growing):** 1 litre/ha in 1,000 litres of water, repeated 3-8 times at 7-14 day intervals. Applications should commence as soon as there is sufficient leaf cover (preventative treatment) or as soon as first symptoms are seen (curative treatment).

**Sugar Beet:** 1 litre/ha in 1,000 litres of water, repeated 3-8 times at 7-14 day intervals. Applications should commence as soon as there is sufficient leaf cover (preventative treatment) or as soon as first symptoms are seen (curative treatment).

**Turf:** 1 l/ha as soon as growth commences in spring and/or following identification of need by analysis. Repeat sprays at 10 to 14 day intervals as necessary. Water rate: 200 l/ha.

## HYDROPONICS

The rate will vary according to the concentration of iron requirement in the final solution. Mixing 10 mls of product with 1000 litres of water will give a solution containing 1 ppm iron.

For further advice and product information please scan the QR code using your smart phone.

UK



ROI



## DIRECTIONS FOR USE

### ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

### TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.

### PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the MSDS for further information, available at [www.yara.com](http://www.yara.com).

### STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

### CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

