



YaraVita™

Zintrac™ 700

NOTICE

Before using this product, read the entire label

Zintrac™ 700

EC FERTILISER

Inorganic foliar fertiliser

Zinc-based fertiliser

Guaranteed analysis:

70% w/v = **700g/l zinc (Zn)** [40% w/w] total as oxide

pH:9

Batch number: see pack for details.

Manufacture date: See pack for details.

Validity date: 2 years from date of manufacture.

Distributed by: **Yara East Africa Ltd.**

P.O. Box 50949, City Square 00200, Nairobi, Kenya.

Tel: +254 0724 255 370 www.yara.co.ke

Distributed by: **Yara Tanzania Ltd.**

P.O.Box 40230 Dar es Salaam, Tanzania.

Tel: +255 737 22 5802/3

Manufactured in the UK by Yara UK Ltd., Pocklington, York YO42 1DN, U.K.

Tel: 00 44 1759 302545

5 litres

Net Weight
8.67Kg

Warning

H410: Very toxic to aquatic life with long lasting effects.

P273: Avoid release to the environment.

P391: Collect spillage.



YaraVita™ Zintrac™ 700

APPLICATION RATES AND TIMINGS

Avocado: 2 l/ha during spring flush, then during summer flush and again during autumn flush (up to 60 days before harvest).

Water rate: minimum 30 l/ha.

Bananas: 1 l/ha. Spray as required. Repeat applications may be necessary where low or marginal levels of zinc exist.

Water rate: minimum 30 l/ha.

Brassicas (field grown): 1 to 2 l/ha at the 4 to 9 leaf stage.

Water rate: minimum 50 l/ha.

Carrot: 1 l/ha when crop is 15 cm tall. For moderate to severe deficiency repeat applications at 10 to 14 day intervals.

Water rate: minimum 200 l/ha.

Cereals: 1 l/ha from 2 leaf stage to first node detectable (Zadok's G.S 12 to 31), repeated once after 10 to 14 days in case of moderate to severe deficiency. Water rate: minimum 50 l/ha.

Coffee: 1 l/ha applied prior to flushes of growth and also post-harvest. Water rate: 200 to 500 l/ha.

Cotton: 1 l/ha 3 to 4 weeks after emergence. Repeat as required with 10 to 14 days between treatments.

Water rate: minimum 50 l/ha.

Eucalyptus: 2 l/ha applied during spring or autumn flush.

Water rate: minimum 20 l/ha.

Garlic: 1 l/ha when sufficient leaf area to intercept spray.

Water rate: minimum 50 l/ha.

Groundnuts/Peanuts: 1 l/ha at the 4 to 6 leaf stage, or sooner when deficiency is visible. Repeat 10 to 14 days later if necessary. Water rate: minimum 200 l/ha.

Maize, Sorghum: 1 l/ha from 3 to 8 leaf stage. For severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals. Water rate: minimum 30 l/ha.

Peas (field grown): 1 l/ha when crop is 5 to 15 cm tall.

Water rate: minimum 50 l/ha.

Pepper (field grown): 1 l/ha applied at the 4 to 6 leaf stage.

Water rate: minimum 500 l/ha.

Pineapple: 3 to 4 applications of 0.5 l/ha at 4 to 6 week intervals from start of leaf growth in early spring.

Water rate: minimum 200 l/ha.

Rice: 1 l/ha applied at start of tillering and again at panicle initiation. Water rate: minimum 50 l/ha.

Sugar Cane: 1 l/ha when cane is between 30 and 120 cm tall. Repeat applications may be necessary at 10 to 14 day intervals. Water rate: minimum 30 l/ha.

Sunflower: 1 l/ha from the 2 pairs of leaves stage.

Water rate: minimum 50 l/ha.

Tea: 1 l/ha applied during plucking season. Repeat at approximately 1 month intervals. Spray 2 to 6 applications.

Water rate: 200 to 500 l/ha. Alternatively: 0.5 to 1 l/ha during the plucking season. Two to six sprays repeated at monthly intervals. Water rate: 20 to 50 l/ha.

Tobacco: 1 l/ha at 3 to 4 leaf stage. Repeat 10 days later if necessary. Water rate: minimum 30 l/ha.

Vegetables (Beans, Onions, Potatoes, Tomatoes (field grown)): 1 l/ha as soon as there is sufficient foliage to intercept a spray. Repeat 10 to 14 days later if necessary.

Water rate: 200 l/ha.

Protected Crops: 0.05 litres per 100 litres water maximum concentration. Water rate: 1000 l/ha maximum. Refer to equivalent field grown crop for application timing.

For further crop recommendations contact your local Yara representative.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the SDS for further information, available at www.yara.com.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.