



YaraVita™
Magphos™ K

NOTICE

Before using this product, read the entire label

Magphos™ K

EC FERTILISER

PK (Mg) fertiliser solution

Guaranteed analysis - soluble in water:

29.7% w/w = **440 g/l Phosphate (P₂O₅)**

5% = w/w **74 g/l Potassium Oxide (K₂O)**

4.1% = w/w **60 g/l Magnesium (Mg)**

Manufactured by: Yara UK Ltd., Pocklington, York,
YO42 1DN, UK

Tel: 01759 302545

Distributed by: **Yara International (M) Sdn Bhd**,
Lot 3.02, Level 3, 1 First Avenue, Bandar Utama
47800 Petaling Jaya, Selangor Darul Ehsan,
Malaysia
Tel: 603 76138010

1 litre

Net weight
1.47 kg

APPLICATION RATES AND TIMINGS

Beans, Peas: 1-2 applications of 5 l/ha when crop is approximately 10-15 cm tall. Allow 10-14 days between treatments. Water rate: 200 l/ha.

Broccoli, Brussel Sprouts, Cabbage, Calabrese, Cauliflower:

4 l/ha applied soon after transplanting. Water rate: 200 l/ha.

Carrot: 3 to 5 l/ha when crop is 15 cm tall. Repeat if necessary at 10 to 14 day intervals. Water rate: 200 l/ha.

Cereals: 5 l/ha at tillering. Repeat at 10-14 day intervals if necessary. Also apply from ear emergence until the end of flowering (Zadok's G.S. 51 to 69). Water rate: 200 l/ha.

Daffodil, Lily, Tulip (Field Grown): 5 l/ha applied 7 to 21 days after the flowers have been removed, and repeated again 10 to 14 days later.

Water rate: 200 l/ha. Do not apply to crops grown under glass or plastic. Do not apply to the crop in flower.

Grass (Armenity): 5 to 10 l/ha as soon as growth commences in spring and/or following identification of need by analysis. Repeat sprays at 10 to 14 day intervals as necessary. Water rate: 200 l/ha

Leek: 5 l/ha applied from two weeks after transplanting, or in the case of direct sown crops when the crop is 15 cm tall. Repeat applications may be necessary at 10 to 14 day intervals. Water rate: 200 l/ha.

Maize: 5 l/ha at 4 to 8 leaf stage. Repeat at 10 to 14 day intervals if necessary. Water rate: 200 l/ha.

Oilseed Rape: For a single application, 5 l/ha at onset of stem extension. For moderate deficiency, 5 l/ha at 4 to 6 leaf stage and at onset of stem extension. An extra application can be made 10 to 14 days later for severe deficiency. Avoid flowering. Water rate: 200 l/ha

Onion: 5 l/ha. Apply when the foliage is 15 cm tall, with a second application 10 to 14 days later if necessary. Also, 1 to 2 applications of 5 l/ha during bulb filling, with a 10 to 14 day interval between sprays. Water rate: 200 l/ha.

Parsnip: 5 l/ha when crop is 15 cm tall. For moderate to severe deficiency repeat applications at 10 to 14 day intervals. (3 applications maximum). Water rate: 200 l/ha.

Plum: 2 to 5 applications of 3 to 5 l/ha from petal fall. Repeat applications at 10 to 14 day intervals. Water rate: 500 to 1000 l/ha.

Potatoes: To increase tuber number, 10 l/ha at tuber initiation (when 50% of the tip swellings are twice the diameter of the rest of the stolon). To increase tuber size a minimum of 2 applications of 5 l/ha or 1 to 2 applications of 10 l/ha during tuber bulking (as soon as first formed tubers are 10 mm in diameter). Allow 10-14 days between applications. Water rate: 200 l/ha

Sugar Beet: 5 l/ha at 4-6 leaf stage. Repeat applications may be made at 10-14 day intervals. Water rate: 200 l/ha.

Turf: 5 to 10 l/ha as soon as growth commences in spring and/or following identification of need by analysis. Repeat sprays at 10 to 14 day intervals as necessary. Water rate: 200 l/ha.

PROTECTED CROPS

0.3 litres per 100 litres water maximum concentration. Water rate: 1000 l/ha maximum. Refer to equivalent field grown crop for application timing

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS
Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 40°C) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



Danger

H290: May be corrosive to metals.

H314: Causes severe skin burns and eye damage.

P260: Do not breathe gas or vapour.

P280: Wear protective gloves/clothing and eye/face protection.

P234: Keep only in original container.

P305+P351+P338+P310: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician.

P303+P361+P353: IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water.