



YaraVita™

Molytrac™
250



1 litre

Net Weight
1.61kg

NOTICE

Before using this product, read the entire label

GROUP 2 FERTILIZER

REGISTRATION No. B3709 Act 36 of 1947

Molybdenum fertilizer in solution for application to deficient crops

Guaranteed analysis - soluble in water:

155g/kg = 250g/l molybdenum (Mo) total

Typical density 1.61kg/litre at 20°C

Manufactured by Yara UK Ltd., Pocklington, York YO42 1DN, U.K.
Registered by Yara Africa Fertilizers (PTY) Ltd.,
Reg.nr. 2011/003434/07
The Pivot @ Montecasino, No. 1 Montecasino Boulevard,
Fourways, Johannesburg 2191.

RECOMMENDED APPLICATION RATES AND TIMINGS

Brassicas: 250 to 300 ml/ha from 4 to 6 leaf stage. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 days intervals. Water rate: 50 to 500 l/ha.

Citrus: 0.25 l/ha from 2/3 of new leaf development in the spring. For moderate to severe deficiency repeat application 10 to 14 days later. Also, 0.25 l/ha applied during autumn flush. Repeat 10 to 14 days later if necessary. Water rate: 500 l/ha.

Cucurbits (field grown): 250 ml/ha one to two weeks after germination and at the 4 leaf stage. Water rate: 200 l/ha

Grapevines: 0.5 l/ha applied before flowering. Repeat at fruit set, bunch closure and early veraison if necessary. Water rate: 200 l/ha.

Legumes, Pulses: 250 to 500 ml/ha when plants are 5 to 15 cm tall. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals from the first application. Water rate: 30 to 200 l/ha. Seed Dressing: 3 litres per tonne.

Lucerne: 250 to 500 ml/ha every 2 to 3 cuts. Water rate: 50 to 200 l/ha. Seed Dressing: 2-3 litres per tonne.

Maize: 0.25 to 0.5 l/ha at the 6 to 8 leaf stage. Water rate: 50 to 200 l/ha. Seed Dressing: 2-4 litres per tonne.

Peas: 250 to 500 ml/ha when plants are 5 to 15 cm tall. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals from the first application. Water rate: 50 to 200 l/ha. Seed Dressing: 1-2 litres per tonne.

Pineapple: 0.25 l/ha. Apply from start of new leaf growth in spring. Regular applications at 10 to 14 day intervals may be necessary where low or marginal levels of molybdenum exist. N.B. Do not apply during flowering. Water rate: 200 l/ha.

Potatoes: 0.5 l/ha between 100% emergence and tuber initiation. Water rate: 50 to 200 l/ha.

Soya Bean: 0.25 to 0.5 l/ha when crop is 5 to 15 cm tall, repeated at 10 to 14 day intervals if necessary. Water rate: 30 to 200 l/ha. Seed Dressing: 1-2 litres per tonne.

Sunflower: 0.25 to 0.5 l/ha from the 4 leaf to flower bud stages. Water rate: 50 to 200 l/ha. Seed Dressing: 2-4 litres per tonne.

Tomato (field grown): 250 ml/ha when crop is 15 cm tall. Water rate: 50 to 500 l/ha.

For further crop recommendations contact your Yara representative.



DIRECTION FOR USE - SEED TREATMENT ALWAYS FOLLOW THESE LABEL INSTRUCTIONS.

Mix the product thoroughly before use and continue mixing during application. Care should be taken to calibrate the machine accurately, to obtain an evenly applied correct dosage. Product application to most seed is enhanced by dilution with water; typically, addition of 25-35% water based on the recommended product rate will achieve good coverage, but this can be varied according to the moisture content of the grain and the equipment used. This product can be used as part of a typical seed treatment slurry in combination with compatible agrochemicals. In such cases this product can replace part of the water used to prepare the slurry to maintain the overall volume of slurry applied.

DIRECTIONS FOR USE - FOLIAR ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the MSDS for further information, available at www.yara.com.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.