



YaraVita™
Frutrel™

NOTICE

Before using this product, read the entire label

Frutrel™

EC FERTILISER

NP (CaO;MgO) fertiliser suspension with boron (B) and zinc (Zn)

Guaranteed analysis:

4.3% w/w = **69 g/l Nitrogen (N)** total

[4,3% ureic nitrogen]

15% w/w = **240 g/l Phosphorus Pentoxide (P₂O₅)**

[9.2% soluble in neutral ammonium citrate]

12.5% w/w = **200 g/l Calcium (Ca)** total

3.8% w/w = **60 g/l Magnesium (Mg)** total

1.3% w/w = **20 g/l Boron (B)** total

2.5% w/w = **40 g/l Zinc (Zn)** total

Contains extract of *Ascophyllum Nodosum*

Manufactured by: Yara UK Ltd., Pocklington, York,
YO42 1DN, UK

Tel: 01759 302545

Distributed by: **Yara International (M) Sdn Bhd,**

Lot 3.02, Level 3, 1 First Avenue, Bandar Utama

47800 Petaling Jaya, Selangor Darul Ehsan,

Malaysia

Tel: 603 76138010

1 litre

Net weight
1.59 kg

APPLICATION RATES AND TIMINGS

Apple: 5 to 10 l/ha at bud burst. Repeat at petal fall. For premature leaf fall (e.g. Golden Delicious): two to five applications of 5 to 10 l/ha from fruit set onwards, at 10 to 14 day intervals. Do not apply within one month of harvest. Water rate: 200 l/ha.

Cherry: Two applications of 5 to 10 l/ha between bud burst and start of flowering. Also, two to five applications of 5 to 10 l/ha from fruit set onwards, at 7 to 14 day intervals. Do not apply within one month of harvest. Water rate: 200 l/ha to 2000 l/ha.

Grapevines: Three applications of 2.5 to 5 l/ha at stages flower truss visible, flower buds separated, start of flowering and cap fall/fruit set. Water rate: 200 l/ha.

Pears: 5 to 10 l/ha at bud burst. Also, two to five applications of 5 to 10 l/ha from fruit set onwards, at 10 to 14 day intervals. Do not apply within one month of harvest. Water rate: 200 l/ha.

Plum: Two applications of 5 to 10 l/ha between bud burst and start of flowering. Also, two to five applications of 5 to 10 l/ha from fruit set onwards, at 10 to 14 day intervals. Do not apply within one month of harvest. Water rate: 200 to 2000 l/ha.

Strawberry (Field Grown): 5 l/ha at the start of new season leaf growth. Repeat twice at 10 to 14 day intervals. Water rate: 200 l/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS
Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 40°C) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



Warning
Contains Boric acid, zinc salt
H361: Suspected of damaging fertility. Suspected of damaging the unborn child.
H412: Harmful to aquatic life with long lasting effects.
P202: Do not handle until all safety precautions have been read and understood.
P280: Wear protective gloves/clothing and eye/face protection.
P273: Avoid release to the environment.
P308+P313: IF exposed or concerned: Get medical attention.
P501: Dispose of contents and container according to local regulations.