



YaraVita™

Stopit™



10 litres

Net Weight  
13.27kg

#### NOTICE

Before using this product, read the entire label  
Irritating to eyes. Avoid contact with skin. Do not breathe spray.

# Stopit™

GROUP 2 FERTILIZER  
REGISTRATION No. K3625 Act 36 of 1947

**Calcium chloride solution**

Guaranteed analysis:

**121g/kg = 160g/l calcium (Ca)** soluble in water

Typical density 1.327kg/litre at 20°C

Manufactured by Yara UK Ltd., Pocklington, York YO42 1DN, U.K.  
Registered by Yara Africa Fertilizers (PTY) Ltd.,  
Reg.nr. 2011/003434/07  
The Pivot @ Montecasino, No. 1 Montecasino Boulevard, Fourways,  
Johannesburg 2191.

#### Warning

Contains: calcium chloride

H319: Causes serious eye irritation.  
P280-a: Wear eye protection.  
P264-a: Wash hands thoroughly after handling.  
P305+p351+p338+p337+p313: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical attention.



# YaraVita™ Stoptit™

## RECOMMENDED APPLICATION RATES AND TIMINGS

**Apples:** 8 to 10 l/ha repeated at 7 to 14 day intervals from end of flowering and timed so that the final application is made one week before harvest. Water rate: 500 l/ha minimum.

**Avocado:** Apply 250 to 500 mls/100 l water from early fruiting at 7 to 10 day intervals. Based on a maximum TRV of 2,000 l/ha.

**Carrots:** 2 to 3 applications of up to 5 l/ha at 10 to 15 cm stage or when sufficient leaf area to intercept spray. Allow 10 to 14 day intervals between applications. Water rate: 200 l/ha.

**Citrus:** 3 to 5 applications of 10 - 20 l/ha applied at 10 to 14 day intervals commencing when fruit reaches 40 to 50 mm size. Water rate: minimum 1000 l/ha. Alternative, 3 applications of up to 300 mls/100 l applied during the period from fruit expansion to harvest. Allow 10 to 14 day intervals between applications. Based on a maximum TRV of 4,000 l/ha.

**Dry Beans:** 5 l/ha applied at the 4-6 leaf stage (10-15 cm tall). Repeat applications can be made at 10-14 day intervals. Water rate: 200 l/ha.

**Grapevines:** A minimum of three applications, 7-10 l/ha, starting at 4mm berry size and repeating at 10-14 days intervals; treat again at berry colouring. Water rate: 500-1,000 l/ha.

**Groundnuts/Peanuts:** Up to 5 l/ha at start of flowering and again during pod development. Water rate: 50-200 l/ha.

**Lettuce (Field Grown):** 2 to 3 applications of up to 5 l/ha commencing 10-14 days after transplanting or emergence, with 10-14 days between applications. Water rate: 50-500 l/ha.

**Mango:** 3 to 5 applications at a rate of up to 125 mls/100 litres water. Applied from early fruiting to fruit swelling, with 7-10 day intervals between applications. Based on a typical TRV of 2,500 l/ha and a maximum TRV of 4,000 l/ha. Apply sufficient water to provide good cover but avoiding run-off.

**Onions:** Up to 5 l/ha at bulb swelling. Repeat applications at 10-14 day intervals as necessary. Water rate: 75-200 l/ha.

**Pears:** 8 to 10 l/ha repeated at 7 to 14 day intervals from end of flowering and timed so that the final application is made one week before harvest. Water rate: 500 l/ha minimum. Alternatively, 250 mls / 100 l repeated at 7 to 14 day intervals from end of flowering and timed so that the final application is made one week before harvest, based on a typical TRV of 2500 l/ha and a maximum TRV of 4000 l/ha.

**Potatoes:** 2 to 3 applications of up to 5 l/ha commencing at tuber initiation (when 50% of the tip swellings are twice the diameter of the rest of the stolon) with 10-14 day intervals between applications. Water rate: 200 l/ha.

**Protected Crops:** 0.5 litres per 100 litres water maximum concentration. Water rate: 1000 l/ha maximum. Refer to equivalent field grown crop for application timing.

**Stone Fruit (Apricots, Cherries, Nectarines, Peaches, Plums):** 5 to 10 l/ha. Two applications should be made at 4 and 2 weeks before harvest. Water rate: 500 l/ha minimum. Alternatively, up to 250 ml/100 litres. Two applications should be made at 4 and 2 weeks before harvest. Based on a typical TRV of 2,500 l/ha and a maximum TRV of 4,000 l/ha. Apply sufficient water to provide good cover but avoiding run-off.

**Tomato (Field Grown):** Spray very young fruits directly, using up to 0.4 litres of 'Stoptit' per 100 litres of water. Maximum water rate: 1,000 l/ha.

**Water Melon (Field Grown):** 2 to 3 applications of 5 l/ha commencing at fruit set with 7 day intervals between applications. Water rate: 200 l/ha.

**For further crop recommendations contact your Yara representative.**

## DIRECTIONS FOR USE

### ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

## TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.

## PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the MSDS for further information, available at [www.yara.com](http://www.yara.com).

## STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

## CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

