

YaraVita

Power Boost™

10 litre

14.43Kg

NOTICE

Before using this product, read the entire label

Power Boost™

FC FFRTII ISFR

Inorganic foliar fertiliser

Mixture of micro-nutrients boron (B) and zinc (Zn) with Nitrogen (N) and Magnesium (Mg)

Guaranteed analysis - Soluble in water:

4.8% = **69g/l** nitrogen (N),

4.9% = 70% = 40g/l magnesium (Mg)

[8.13% magnesium oxide (MgO)]

4.9% = 70g/l boron (B)

9.9% = 140g/l zinc (Zn)

nH:9.5

Manufacture date: See top of pack.

Expiry date: 2 years from date of manufacture.

Distributed by: Yara East Africa Ltd.

P.O. Box 50949, City Square 00200, Nairobi, Kenya.

Tel: +254 0724 255 370

www.vara.co.ke

Distributed by: Yara Tanzania Ltd.

P.O.Box 40230 Dar es Salaam, Tanzania.

Tel: +255 737 22 5802/3

Manufactured in the UK by Yara UK Ltd., Pocklington, York YO42

1DN, U.K. Tel: 00 44 1759 302545

WARNING

Hazardous ingredients: Boric acid, zinc

H361: Suspected of damaging fertility or the unborn child. H400: Very toxic to aquatic life. H411: Toxic to aquatic life with long lasting effects. P202: Do not handle until all safety precautions have been read and understood. P280: Wear protective gloves/clothing and eve/face protection, P273; Avoid release to the environment.

P308+P313: IF exposed or concerned: Get medical attention, P391: Collect spillage. P501: Dispose of contents and container according to local

regulations.

Net Weight

ENVIRONMENTALLY HAZARDOUS SUBSTANCE. LIOUID, N.O.S. (Boric acid, zinc salt)



WLKEYPD4L00 V.11.24

YaraVita™ Power Boost™

APPLICATION RATES AND TIMINGS

Coffee: 1 l/acre at the start of shoot growth. Repeat if necessary after fruit set. Water rate: 200 l/acre.

Avocado: 800mll/acre during spring and summer flushes. Water rate: 400 l/acre.

Tea: 400 ml/acre applied during plucking season. Repeat at approximately 1 month intervals. Spray 2 to 6

applications. Water rate: 200 l/acre.

Kahawa: lita 1 kwa ekari wakati wa machipukizi mapya. Rudia wakati wa uwekaji matunda kama kuna ulazima. Tumia maji kiasi cha lita 200 kwa ekari.

Parachichi: Miliita 800 kwa ekari wakati wa machipukizi mapya. Kiasi cha maji ni lita 400 kwa ekari.

Chai: Mililita 400 kwa ekari mara baada ya kuvuna. Rudia baada ya mwezi mmoja. Nyunyizia mara 2 hadi 6

kwa msimu. Kiasi cha maji lita 200 kwa ekari.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the SDS for further information, available at www.yara.com.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.