



YaraAmplix™

FLOSTREL™



10 litres
Net weight
13.5 kg

NOTICE

Before using this product, read the entire label

FLOSTREL™

UK FERTILISER

Compound fluid NP fertiliser with boron.

Content % w/w:

5.2% (= 71 g/l) Total Nitrogen (N)

7.3% (= 100 g/l) Total Phosphorus Pentoxide (P₂O₅)

[7.3% Phosphorus Pentoxide (P₂O₅) water soluble]

7.3% (= 100 g/l) Boron (B) as ethanalamine, water soluble.

INFORMATION FOR ROI MARKET

PFC 1(C)(I)(b)(i) Straight liquid inorganic macronutrient fertiliser

P 7.3 with Boron

Liquid Solution

Content % w/w:

7.3% (= 100 g/l) Total Phosphorus Pentoxide (P₂O₅)

[7.3% Phosphorus Pentoxide (P₂O₅) water soluble]

7.3% (= 100 g/l) Boron (B) as ethanalamine, water soluble

Also contains:

5.2% (= 71 g/l) Total Nitrogen (N)

Ingredients: Orthoboric acid, compound with 2-aminoethanol CAS n° 26038-87-9 (CMC 1: Virgin material substances and mixtures), Monoethanolamine CAS n° 141-43-5 (CMC 1), orthophosphoric acid CAS n° 7664-38-2 (CMC 1).

To be used only where there is a recognised need.

Do not exceed the appropriate dose rates

Batch Number/ Manufacture Date: See Top of Pack

Produced and Distributed by: **Yara UK Limited.**

Pocklington Industrial Estate, Pocklington, York. YO42 1DN

Tel: 01759 302545

www.yara.co.uk

ROI: Imported by: **Yara UK Limited.**

The Old Rectory, Cork Road, Carrigaline, P43 W921. County Cork.

Tel: +44 1759 302 545

www.yara.ie

YaraAmplix™ FLOSTREL™

APPLICATION RATES AND TIMINGS

Barley: Apply at 1 l/ha from the 4 leaf stage to pseudo stem erection, (Zadoks Growth Stage 14 to 30). Apply at 0.5 l/ha from first node detectable to the end of booting, (Zadoks G.S. 31 to 49).

Water rate: 100 to 200 l/ha.

Beans: 2 l/ha at a growth height of 20-40cm (BBCH 51-55) and repeated at a growth height of approximately 100cm (beginning of flowering, BBCH 61-65). Water volume: 200 l/ha.

Lupin: 2 l/ha at a growth height of 20-40cm (BBCH 51-55) and repeated at a growth height of approximately 100cm (beginning of flowering, BBCH 61-65). Water volume: 200 l/ha

Maize: 2 l/ha at 3 to 8 leaf stage.

Water volume: 200 l/ha.

Oilseed rape / canola: Apply 1-2 L/ha at the 4 to 9 leaf stage and again at the onset of the rapid stem extension phase up to the start of flowering.

Water volume 100-200L/ha.

Peas: 2 l/ha at growth height 20-30 cm (BBCH 51-55) and repeated at the beginning of flowering (BBCH 61-65). Water volume: 200 l/ha.

Potatoes: 2-3 applications of 1 to 2 l/ha each, 7 to 14 days after full emergence. Repeat if necessary, after 10 - 14 days. Water volume: 200 l/ha

Soya bean: 2 l/ha at a growth height of 20-40cm (BBCH 51-55) and repeated at a growth height of approximately 100cm (beginning of flowering, BBCH 61-65). Water volume: 200 l/ha.

Sugar beet: 3 l/ha from the 4 to 6 leaf stage.

Repeat after 10 to 14 days if necessary.

Water volume: 200 l/ha.

Wheat: Apply at 1 l/ha from the 4 leaf stage to pseudo stem erection, (Zadoks Growth Stage 14 to 30). Apply at 0.5 l/ha from first node detectable to the end of booting, (Zadoks G.S. 31 to 49).

Water rate: 100 to 200 l/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the SDS for further information, available at www.yara.com.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

TANK MIXING/CO-APPLICATION



Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit Tankmix for important information. Alternatively, contact Yara for specific advice.

