YaraVita Amazinc™

RECOMMENDED APPLICATION RATES AND TIMINGS

Almond, Hazelnuts, Pecans, Walnuts: 1 to 2 l/ha; From leaf canopy development through hull split at 14-21 day interval as necessary. Also post-harvest before onset of leaf senescence. Avoid applications during flowering and fruit set. Water rate: 500 l/ha.

Apple: Three applications of 1 l/ha at: green bud; early bud development (e.g. 20 mm diameter); and post-harvest but before leaf senescence. Avoid applications during flowering. Water rate: 500 l/ha minimum.

Cereals: 1 to 2 I/ha from 3 leaf stage to first node detectable (Zadok's G.S. 13 to 31). For moderate to severe deficiency repeat after 10 to 14 days. Water rate: 50 to 200 I/ha. Seed dressing: 3 to 6 I/tonne

Citrus: 2 to 3 l/ha during spring and autumn flushes. Water rate: 500 to 1000 l/ha

Grapevines: 1 l/ha applied at flower buds visible and again at flower buds separated and at fruit set. Water rate: 200 to 1000 l/ha.

Nuts (deciduous): 1 to 2 l/ha at bud break. Repeat 15 days later if necessary. Water rate: 500 to 1000 l/ha.

Onion: 1 to 2 I/ha two weeks after transplanting, or in the case of direct sown crops, when the crop is 15 cm tall. Repeat applications at 10 to 14 day intervals. Water rate: 200 I/ha.

Pears: 1 to 2 l/ha at bud burst. Avoid flowering. Apply 1 to 3 l/ha after harvest but before leaf senescence. Water rate: 500 to 1500 l/ha.

Peas: 1 l/ha at 10 to 15 cm (4 to 6 leaf) stage. Repeat as necessary at 7 to 14 day intervals. Avoid application during flowering. Water rate: 30 to 100 l/ha.

Stone Fruit: 1 to 2 l/ha during fruit development. Repeat if necessary at 10 to 14 day intervals up to one month before harvest. Water rate: 500 l/ha

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes (temperatures exceeding 40°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

Warning

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

For further crop recommendations visit Product Information at www.yara.com.au TOLL FREE number 1800 684 266, Emergency tel: 02 8014 4558

