YaraVita BioMaris™

APPLICATION RATES AND TIMINGS

Apple, Apricot, Avocado, Cherry, Citrus: 1-2 l/ha. Apply 4-5 treatments covering critical growth stages such as vegetative growth stage, petal fall and early fruit sizing. Application rates should be adjusted based on plant size and leaf area. Use enough water for good spray coverage. Maximum concentration for foliar applications is 500 ml per 100 litres of water.

Broccoli: 1-2 I/ha. Apply 3-4 treatments at 10 to 15 days intervals as necessary. Start applications 10 days after seedling emergence or from 3-4 leaf stage. Water volume: 200 I/ha minimum

Carrot: 1-2 l/ha. Apply 2-3 treatments at 10-15 days interval. Commence the applications when roots or tubers start to

develop. Water volume: 200 l/ha minimum

Cereals: 1-2 I/ha. Apply 3 treatments. Commence the application at 5 leaf stage and cover critical growth stages such as tillering, stem extension, or flag leaf stage. Water volume: 200 I/ha minimum

Grapes (wine): 1-2 I/ha. Apply 4-5 treatments covering critical growth stages such as vegetative growth stage, petal fall and early fruit sizing. Application rates should be adjusted based on plant size and leaf area. Use enough water for good spray coverage. Water volume: 200 I/ha minimum

Kiwifruit: 1-2 I/ha. Apply 4-5 treatments covering critical growth stages such as vegetative growth stage, petal fall and early fruit sizing. Application rates should be adjusted based on plant size and leaf area. Use enough water for good spray coverage. Maximum concentration for foliar applications is 500 ml per 100 litres of water.

Lettuce: 1-2 I/ha. Apply 3-4 treatments at 10 to 15 days intervals as necessary. Start applications 10 days after seedling emergence or from 3-4 leaf stage. Use enough water for good spray coverage. Maximum concentration for foliar applications is 500 ml per 100 litres of water.

Onion: 1-2 l/ha. Apply 3-4 treatments at 10 to 15 days intervals as necessary. Start applications 10 days after seedling emergence or from 3-4 leaf stage. Water volume: 200 l/ha minimum

Potatoes:1-2 I/ha. Apply 2-3 treatments at 10-15 days interval. Commence the applications when roots or tubers start to develop. Water volume: 200 I/ha minimum

Squash: 1-2 l/ha. Apply 4-5 treatments covering critical growth stages such as vegetative growth stage, petal fall and early fruit sizing. Application rates should be adjusted based on plant size and leaf area. Use enough water for good spray coverage. Maximum concentration for foliar applications is 500 ml per 100 litres of water.

Strawberry: 1-2 I/ha. Apply 4-5 treatments covering critical growth stages such as vegetative growth stage, petal fall and early fruit sizing. Application rates should be adjusted based on plant size and leaf area. Use enough water for good spray coverage. Maximum concentration for foliar applications is 500 ml per 100 litres of water.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes (temperatures exceeding 40°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

