



**YaraVita™**

**Bortrac®150**



**10 litres**  
**Net weight**  
**13.53 kg**

#### NOTICE

Before using this product, read the entire label

# Bortrac®150

#### UK FERTILISER

Boron ethanolamine

Content % w/w:

**11.5% Total Boron (B)** as ethanolamine,  
water soluble

Also Contains % w/w:

**4.8% Total Nitrogen (N)**  
[4.3% Ammoniacal Nitrogen, 0.5% Urea Nitrogen]

To be used only where there is a recognised need.  
Do not exceed the application rate

#### INFORMATION FOR ROI MARKET

**PFC 1(C)(II)(a) Straight Inorganic micronutrient fertilizer**

Content % w/w:

**11.5% Total Boron (B)** as ethanolamine,  
water soluble

Also Contains % w/w:

**4.8% Total Nitrogen (N)**  
[4.3% Ammoniacal Nitrogen, 0.5% Urea Nitrogen]

**Ingredients:** orthoboric acid, compound with 2-aminoethanol CAS<sup>o</sup>  
26038-87-9 (CMC 1: Virgin material substances and mixtures)

To be used only where there is a recognised need.  
Do not exceed the application rate.

This fertiliser contains urea, which can release ammonia and have an impact on air quality. Depending on local conditions, appropriate remediation measures must be taken.

**Batch Number:** See Pack      **Manufacture Date:** See Pack

Distributed by: **Yara UK Limited**. Harvest House, Europarc,  
Grimsby, N E Lincolnshire, DN37 9TZ.  
Tel:+44 1472 889 250    [www.yara.co.uk](http://www.yara.co.uk)

**ROI:** Imported by: **Yara UK Ltd**. The Old Rectory, Cork Road,  
Carrigaline, P43 W921. County Cork.  
Tel: +44 1472 889 250    [www.yara.ie](http://www.yara.ie)

Produced by: **Yara UK Ltd.**, Manor Place, Pocklington,  
York, YO42 1DN, UK.

# YaraVita™ Bortrac®150

For professional use only.

Please contact Yara for more specific recommendations.

## APPLICATION RATES AND TIMINGS

### FOLIAR APPLICATION

**Apples, Pears:** 1 to 2 l/ha at pink bud, start of flowering and again at petal fall. 2 l/ha after harvest but before leaf senescence. Water rate: 500-1,000 l/ha.

**Beans, Peas:** 2 l/ha at 10-15 cm stage. For moderate to severe deficiency, a repeat application may be necessary 10-14 days later. Water rate: 200 l/ha.

**Broccoli, Brussel Sprouts, Cabbage, Calabrese,**

**Cauliflower:** 3 l/ha at 4-6 leaf stage with repeat applications at the above rate at 10-14 day intervals for moderate to severe deficiency. Water Rate: 200 l/ha.

**Carrot:** 3 l/ha at 10-15 cm stage or when sufficient foliage to intercept spray. For moderate to severe deficiency, repeat applications at 10 to 14 days later.

Water rate: 200 l/ha.

**Cherry, Plum:** 1-2 l/ha at winter bud and again at pink bud. 2 l/ha after harvest but before leaf senescence.

Water rate: 500-1,000 l/ha.

**Oilseed Rape:** For a single application. 3 l/ha at onset of stem extension. For moderate deficiency, 3 l/ha at 4 to 6 leaf stage and 3 l/ha at onset of stem extension. An extra application can be made between 10 and 14 days later or up to the start of flowering. Water rate: 50 to 200 l/ha.

**Onion:** 1 l/ha as soon as there is sufficient foliage to intercept spray. A second application may be made 10 to 14 days later. Water rate: 200 l/ha.

**Potatoes:** One application of 1 to 2 l/ha applied one week after 100% emergence. A second application may be made 10 to 14 days later. Water rate: 200 l/ha.

**Strawberry (Field Growth):** 2 applications of 1 l/ha commencing at white bud and repeated 10-14 days later. Also, 2 l/ha applied at regrowth (after harvest).

Water rate: 200-500 l/ha.

**Red Beet, Sugar Beet, Swede, Turnip:** 3 l/ha at 4-6 leaf stage. For moderate to severe deficiency, repeat applications should be made at the above rate at 10-14 day intervals. Water rate: 50-200 l/ha.

### PROTECTED CROPS

0.1 litres per 100 litres water maximum concentration. Water rate: 1000 l/ha maximum. Refer to equivalent field grown crop for application timing.

### SOIL APPLICATION

**All Crops:** 5 l/ha pre-planting or pre-emergence in sufficient water to ensure accurate application.

For further advice and product information please scan QR code using your smart phone.

UK



ROI



## DIRECTIONS FOR USE

### ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

### TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.

### PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the MSDS for further information, available at [www.yara.com](http://www.yara.com).

### STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

### CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

