



YaraVita™

Tracel™ BZ



NOTICE

Before using this product, read the entire label

Tracel™ BZ

EC FERTILISER

Inorganic foliar fertiliser

Mixture of micronutrients boron (B), copper (Cu), iron (Fe), manganese (Mn), molybdenum (Mo) and zinc (Zn) with nitrogen (N), potassium (K₂O) and phosphorus (P₂O₅) with magnesium (Mg) and sulphur (S)

Guaranteed analysis:

5% boron (B) as borate, **5% zinc (Zn)** as sulphate/phosphate, **0.1% copper (Cu)** as EDTA, **0.1% iron (Fe)** as citrate, **0.1% manganese (Mn)** as sulphate, **0.1% molybdenum (Mo)** as molybdate, **5% nitrogen (N)** as urea, **7.5% phosphate (P₂O₅)** as zinc/potassium, **5% potash (K₂O)** as chloride/phosphate, **5% magnesium (Mg)** as oxide.

To be used only where these is recognised need

Do not exceed the appropriate application rates

pH in 2.5% solution: 8

Manufacture date: See top of pack

Expiry date: 5 years from date of manufacture.

Distributed by: **Yara East Africa Ltd.**

P.O. Box 50949, City Square 00200, Nairobi, Kenya.

Tel: +254 0724 255 370

www.yara.co.ke

Distributed by: **Yara Tanzania Ltd.**

P.O.Box 40230 Dar es Salaam, Tanzania.

Manufactured in the UK by Yara UK Ltd., Pocklington, York YO42 1DN, U.K.

Tel: 00 44 1759 302545

Danger

Hazardous ingredients: disodium octaborate, zinc sulphate (anhydrous)

H319: Causes serious eye irritation.

H360: May damage fertility or the unborn child. H411: Toxic to aquatic life with long lasting effects. P202: Do not handle until all safety precautions have been read and understood. P280-d: Wear protective gloves/clothing and eye/face protection.

P305+P351+P338: IF IN

EYES: Rinse cautiously with

water for several minutes.

Remove contact lenses, if

present and easy to do. Continue

rinsing. P308+P313: IF exposed or

concerned: Get medical attention.


P391: Collect spillage. P501: Dispose of

contents and container according to

local regulations.

UN3077

ENVIRONMENTALLY HAZARDOUS SUBSTANCE,
SOLID, N.O.S. (trizinc bis(orthophosphate), zinc sulphate)



Net Weight
20Kg

YaraVita™ Tracel™ BZ

RECOMMENDED APPLICATION RATES AND TIMINGS

Coffee: 2 kg/ha applied early in the growing season. Repeat at the same rate at the pre-flowering stage and also post-harvest. Water rate: 500 to 1000 l/ha.

Bananas, Citrus, Mango: 2 kg/ha applied to new leaf growth in the spring and again 10 to 14 days later. Repeat the application after final harvest. N.B. Do not apply during flowering. Water rate: 600 to 1000 l/ha.

Beans, Peas: 2 to 4 kg/ha at the 4 to 6 leaf stage. A second application may be made 10 to 14 days later. Water rate: 200 to 400 l/ha.

Cabbage, Kale: 2 kg/ha at the 4 to 6 leaf stage, with repeat applications at 10 to 14 day intervals. Water rate: 200 to 500 l/ha.

Capsicums (Field Grown), Tomato (Field Grown): 2 kg/ha when the plants are at the 4 to 6 leaf stage. Repeat once or twice at 10 to 14 day intervals. Water rate: 200 to 500 l/ha.

Cereals (Barley, Rice, Wheat): 2 to 4 kg/ha at the 4 to 6 leaf stage. Water rate: 200 to 400 l/ha.

Cocoa / Cacao: 2 kg/ha applied early in the growing season. Repeat at the same rate at the pre-flowering stage and also post-harvest. Water rate: 500 to 1000 l/ha.

Forestry (Conifers, Eucalyptus): Two applications of 2 to 4 kg/ha when there is new season leaf production. Water rate: 200 to 1000 l/ha.

Maize, Sorghum: 2 to 4 kg/ha applied from 4 to 6 leaves. Water rate: 200 to 400 l/ha.

Onion: 1 to 2 kg/ha as soon as there is sufficient foliage to intercept the spray. A second application may be made at the same rate 10 to 14 days later. Water rate: 200 to 500 l/ha.

Pineapple: 2 kg/ha applied to new leaf growth in the spring and again 10 to 14 days later. Repeat the application after final harvest. Water rate: 200 to 1000 l/ha.

Potatoes: 2 kg/ha 7 to 14 days after full emergence. Repeat at tuber initiation and at tuber bulking. Water rate: 200 to 500 l/ha.

Sugar Cane: 2 kg/ha when the cane is between 30 and 90 cm tall. Repeat the application 10 to 14 days later. Water rate: 200 to 500 l/ha.

Sunflower, Oilseed Rape: 2 to 4 kg/ha from 4 to 6 leaves. Consider a second application 10 to 14 days later. N.B. Do not apply during flowering. Water rate: 200 to 400 l/ha.

Tea: 2 kg/ha applied to new leaf growth in the spring and again 10 to 14 days later. Repeat the application after final harvest. N.B. Do not apply during flowering. Water rate: 600 to 1000 l/ha.

For further crop recommendations contact your Yara representative.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS
Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the SDS for further information, available at www.yara.com.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.