



YaraVita™
Safe K™

NOTICE
Before using this product, read the entire label

Safe K™

EC FERTILISER

NK fertiliser solution

Guaranteed analysis - soluble in water:
3.1% w/w = **45 g/l Nitrogen (N)**
34.3% w/w = **500 g/l Potassium Oxide (K₂O)**

Manufactured by: Yara UK Ltd., Pocklington, York,
YO42 1DN, UK
Tel: 01759 302545
Distributed by: **Yara International (M) Sdn Bhd**,
Lot 3.02, Level 3, 1 First Avenue, Bandar Utama
47800 Petaling Jaya, Selangor Darul Ehsan,
Malaysia
Tel: 603 76138010

1 litre

Net weight
1.45 kg

DOSE RATES AND TIMINGS

Apple: 5 l/ha at bud burst. Also, two to three applications of 2 to 3 l/ha at 10 to 14 day intervals starting at petal fall. Water rate: 500 l/ha.

Beans, Peas: 5 l/ha before flowering. Water rate: 30 to 200 l/ha.

Grapevines: 3 to 5 l/ha as soon as there is sufficient new season leaf growth to intercept a spray with up to two repeat applications at 10 to 14 day intervals prior to flowering and/or 3 to 5 l/ha at fruit set, pea-sized berries, first colour softening/one month before harvest. Water rate : 200 to 500 l/ha.

Maize: 5 l/ha at the 4 to 8 leaf stage. Water rate: 30 to 200 l/ha.

Oilseed rape / canola: 5 l/ha at the 4 to 6 leaf stage. Repeat as required for moderate to severe deficiency at 7 to 14 day intervals. Water rate: 200 l/ha.

Potatoes: One application of 10 l/ha or two applications of 5 l/ha during tuber bulking (as soon as first-formed tubers are 10 mm in diameter). Allow 10 to 14 days between applications. Also, 5 l/ha following petiole analysis, during tuber bulking. Water rate: 200 l/ha.

Soya bean: 3 l/ha when the crop is 10 to 15 cm tall. Water rate: 30 to 200 l/ha.

Wheat: 5 l/ha from mid-tillering to 2nd node detectable (Zadoks G.S. 25 to 32). Repeat at 10 to 14 day intervals between these growth stages as necessary. Water rate: 200 l/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS
Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 40°C) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.