



**YaraVita™**  
**Seniphos®**



**10 litres**  
**Net weight**  
**13.12 kg**

# Seniphos®

## NOTICE

Before using this product, read the entire label

## UK FERTILISER

NP (Ca) fertiliser solution 3.1-25 (3)

Guaranteed analysis - soluble in water:

**3.1% (= 41 g/l) Nitrogen (N) total**  
[2% Ammoniacal Nitrogen, 1.1% Nitric Nitrogen],  
**25% (= 328 g/l) Phosphorus Pentoxide (P<sub>2</sub>O<sub>5</sub>)**  
[10.9% (= 143 g/l) Phosphorus (P)]  
**4.2% (= 55 g/l) Calcium Oxide (CaO)**  
[3% (= 40 g/l) Calcium (Ca)]

## INFORMATION FOR ROI MARKET

### PFC 1(C)(I)(b)(ii) Compound liquid Inorganic macronutrient fertilizer

NP (Ca) 3.1- 25 (+4.2)

Liquid Solution

**Ingredients:** Mono potassium phosphate CAS n° 7778-77-0 (CMC 1: Virgin material substances and mixtures), orthophosphoric acid CAS n° 7664-38-2 (CMC 1), calcium bis(dihydrogenphosphate) CAS n° 10031-30-8 (CMC 1), calcium nitrate tetrahydrate CAS n° 13477-34-4 (CMC 1).

Batch Number/ Manufacture Date: See Top of Pack


Produced and Distributed by: **Yara UK Limited**. Pocklington Industrial Estate, Pocklington, York. YO42 1DN Tel: 01759 302545 [www.yara.co.uk](http://www.yara.co.uk)

**ROI:** Imported by: **Yara UK Limited**. The Old Rectory, Cork Road, Carrigaline, P43 W921. County Cork. Tel: +1759 302 545 [www.yara.ie](http://www.yara.ie)

UFI: 4844-5089-G004-BUGH

## Danger

Hazardous ingredients: calcium bis(dihydrogenorthophosphate), phosphoric acid, calcium nitrate tetrahydrate  
H290: May be corrosive to metals. H314: Causes severe skin burns and eye damage. P260: Do not breathe gas or vapour. P280: Wear protective gloves/clothing and eye/face protection. P305+P351+P338+P310: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician. P303+P361+P353: IF ON SKIN (or hair): Take off immediately all contaminated clothing. \*Rinse skin with water. P234: Keep only in original packaging.



**UN 3264**  
CORROSIVE LIQUID, ACIDIC,  
INORGANIC, N.O.S  
(contains phosphoric acid)

# YaraVita™ Seniphos®

## APPLICATION RATES AND TIMINGS

**Apple:** 5-8 applications of 10 l/ha at 10-14 day intervals commencing at petal fall. Water rate: 500 l/ha.

For red coloration: Seniphos at 10 l/ha in 1,000 litres of water has been shown to enhance red colouration. 1 or 2 applications (7 days apart) should be made once the fruit has started to change colour, normally 2 to 3 weeks before harvest.

**Blackcurrant:** 3 applications of 10 l/ha. 50% fruit set, 100% fruit set, and fruit swelling. Water rate: 500 l/ha.

**Broccoli, Brussel Sprouts, Cabbage, Calabrese,**

**Cauliflower:** 2 to 3 applications of 10 l/ha from stem extension/head development at 7 to 14 day intervals. Water rate: 200 l/ha.

**Carrot:** 5 l/ha. 2 to 3 applications when crop is 15 cm tall and at 7 to 10 day intervals. Water rate: 200 l/ha.

**Cereals:** 5 l/ha at tillering. Repeat at 10-14 day intervals if necessary. Water rate: 200 l/ha.

**Cherry, Plum:** 2 to 5 applications of 10 l/ha at 7 to 14 day intervals commencing at petal fall. Water rate: 500 l/ha.

**Hops:** 4 applications of 10 l/ha commencing when crop starts to burr, with 3 further applications at 14 day intervals. Water rate: 500 l/ha.

**Lettuce (Field Grown):** 2 to 3 applications of 5 l/ha commencing 10 to 14 days after transplanting or emergence with 7 to 10 day intervals between applications. Water rate: 200 to 500 l/ha.

**Onion:** 1 to 2 applications of 5 l/ha during bulb filling, with a 10 to 14 day interval between sprays. Water rate: 200 l/ha.

**Pears:** 5 to 8 applications of 10 l/ha at 10 to 14 day intervals commencing at petal fall. Water rate: 500 l/ha.

**Raspberry:** 3 applications of 10 l/ha. Start of flowering, end of flowering, and fruit development. Water rate: 500 l/ha

**Strawberry (Field Grown):** Non-everbearing varieties, 3 applications of 10 l/ha from start of flowering. Repeat applications at 7-10 day intervals. Everbearing varieties, divide a total rate of 30 l/ha into 6 applications of 5 l/ha. Do not apply successive applications at intervals of less than 10-14 days. Water rate: 500 l/ha.

## PROTECTED CROPS

0.5 litres per 100 litres water maximum concentration.

Water rate: 1000 l/ha maximum.

Refer to equivalent field grown crop for application timing.

For further advice and product information please scan the QR code using your smart phone.

UK



ROI



## DIRECTIONS FOR USE

### ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

### TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.

### PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the MSDS for further information, available at [www.yara.com](http://www.yara.com).

### STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

### CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

