



**YaraVita™**  
**Stopit™**



**10 litres**  
**Net weight**  
**13.27 kg**

## NOTICE

Before using this product, read the entire label

# Stopit™

## UK FERTILISER

Calcium chloride solution

Guaranteed analysis:

**16.9% w/w calcium oxide (CaO)** water soluble  
[12% w/w = 160 g/l calcium (Ca)]

## INFORMATION FOR ROI MARKET

**PFC 1(C)(I)(b)(i) Straight liquid Inorganic macronutrient fertilizer**

Ca (16.9)

Liquid Solution

Content % w/w:

**16.9% w/w calcium oxide (CaO)** water soluble  
[12% w/w = 160 g/l calcium (Ca)]

**Ingredients:** Calcium chloride CAS n° 10043-52-4 (CMC 1: Virgin material substances and mixtures).

Batch Number/ Manufacture Date: See Top of Pack

Produced and Distributed by: **Yara UK Limited.**

Pocklington Industrial Estate, Pocklington, York. YO42 1DN

Tel: 01759 302545

[www.yara.co.uk](http://www.yara.co.uk)

**ROI:** Imported by: **Yara UK Limited.** The Old Rectory, Cork Road, Carrigaline, P43 W921. County Cork.

Tel: +44 1472 889 250

[www.yara.ie](http://www.yara.ie)

## Warning

Hazardous ingredients:

calcium chloride. H319:

Causes serious eye irritation.

P280-a: Wear eye protection.

P264-a: Wash hands

thoroughly after handling.

P305+P351+P338+P337+

P313-a: IF IN EYES: Rinse

cautiously with water for

several minutes. Remove

contact lenses, if present and

easy to do. Continue rinsing. If

eye irritation persists: Get

medical attention.



# YaraVita™ Stopit™

## APPLICATION RATES AND TIMINGS

**Apple:** 2 to 5 l/ha applied at bud burst and repeated at 7 to 10 day intervals as necessary to the beginning of flowering. Regular applications of 5 to 10 l/ha at 7 to 14 day intervals from petal fall to one week before harvest. Water rate: 150 to 500 l/ha.

**Broccoli, Brussel Sprouts, Cabbage, Calabrese, Cauliflower:** 2 to 3 applications of 5 l/ha from stem extension/head development to harvest. Allow 10 to 14 days between applications. Water rate: 200 l/ha.

**Carrot:** 2 to 3 applications of 5 l/ha at 10 to 15 cm stage or when sufficient leaf area to intercept spray. Allow 10 to 14 day intervals between applications. Water rate: 200 l/ha.

**Celery:** 5 l/ha early in the season. Repeat at 10 to 14 day intervals as necessary. Water rate: 200 l/ha.

**Cereals:** 5 l/ha from 4 leaves to second node detectable (Zadok's G.S. 14 to 32). Water rate: 200 l/ha

**Cherry, Plum:** 2 applications of 10 l/ha, 4 weeks and 2 weeks before harvest. Water rate: 500 l/ha.

**Leek:** 5 l/ha applied two weeks after transplanting, or in the case of direct sown crops, when the crop is 15 cm tall. Repeat applications may be necessary at 10 to 14 day intervals. Water rate: 200 l/ha.

**Lettuce (Field Grown):** 2 to 3 applications of 5 l/ha commencing 10 to 14 days after transplanting or emergence, with 10 to 14 day intervals between applications. Water rate: 500 l/ha.

**Daffodil, Lily, Tulip (Field Grown):** 5 l/ha applied 7 to 21 days after the flowers have been removed, and repeated again 10 to 14 days later. Water rate: 200 l/ha. Do not apply to crops grown under glass or plastic. Do not apply to the crop in flower.

**Onion:** 5 l/ha at bulb swelling. Repeat applications at 10 to 14 day intervals as necessary. Water rate: 200 l/ha.

**Parsnip:** 5 l/ha when crop is 15 cm tall. For moderate to severe deficiency, repeat applications at 10 to 14 day intervals. (3 applications maximum). Water rate: 200 l/ha.

**Pears:** Regular applications of 5 to 10 l/ha at 7 to 14 day intervals from petal fall to one week before harvest. Water rate: 150 to 500 l/ha.

**Potatoes:** 2 to 3 applications of 5 to 10 l/ha commencing at tuber initiation (when 50% of the tip swellings are twice the diameter of the rest of the stolon) with 10 to 14 day intervals between applications. Water rate: 200 l/ha

**Strawberry (Field Grown):** 3 applications of 10 l/ha from start of flowering. Repeat applications at 7 to 10 day intervals. Water rate: 300-600 l/ha.

## PROTECTED CROPS

0.5 litres per 100 litres water maximum concentration. Water rate: 1000 l/ha maximum. Refer to equivalent field grown crop for application timing.

For further advice and product information please scan the QR code using your smart phone.

UK



ROI



## DIRECTIONS FOR USE

### ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

### TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.

### PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the MSDS for further information, available at [www.yara.com](http://www.yara.com).

### STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

### CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

