

YaraVita Molytrac™ 250

APPLICATION RATES AND TIMINGS

Beans, Peas: 0.25 l/ha when plants are 5 to 15 cm tall. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals from the first application. Water rate: 200 l/ha.

Broccoli, Brussels Sprouts, Cabbage, Calabrese: 0.25 l/ha from 4 to 6 leaf stage. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals. Water rate: 200 l/ha.

Cauliflower: 0.25 l/ha from 4 to 6 leaf stage. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals, or a second application may be made at early head development or buttoning. Water rate: 200 l/ha.

Clover: Apply 0.125 to 0.25 l/ha to new growth in the spring. Water rate: 50 to 200 l/ha. (For Seed Production): Apply 0.25 l/ha just after every cut. Water rate: 50 to 200 l/ha.

Grass: Max Rate of Single Application is 0.5 l/ha; Water rate: 50 l/ha; As soon as growth commences in the spring. Repeat at 7 to 14 day intervals as necessary. Warning: Excessive molybdenum can be harmful to stock. Do not apply fertiliser containing molybdenum to more than 25% of farm annually. Do not graze for 4 weeks after application.

Kale: 0.25 l/ha from 4 to 6 leaf stage. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals. Water rate: 200 l/ha.

Leek: 0.25 l/ha applied two weeks after transplanting, or in the case of direct sown crops, when the crop is 15 cm tall. One or two repeat applications may be necessary at 10 to 14 day intervals. Water rate: 200 l/ha.

Oilseed Rape: For a single application, 0.25 l/ha at onset of stem extension. For moderate deficiency, 0.25 l/ha at 4 to 6 leaf stage and 0.25 l/ha at onset of stem extension. An extra application can be made 10 to 14 days later for a severe deficiency. Avoid flowering. Water rate: 200 l/ha.

Onion: 0.25 l/ha applied as soon as there is sufficient leaf cover to intercept the spray. Repeat two weeks later if necessary. Water rate: 200 l/ha.

Parsnip: 0.25 l/ha 10 to 14 days after transplanting or when crop is 15 cm tall. Water rate: 200 l/ha.

Swede, Turnip: 0.25 l/ha from 4 to 6 leaf stage. For moderate to severe deficiency, one or two repeat applications should be made at the above rate at 10 to 14 day intervals. Water rate: 200 l/ha.

PROTECTED CROPS

0.025 litres per 100 litres water maximum concentration. Water rate: 1000 l/ha maximum.

Refer to equivalent field grown crop for application timing.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS
Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 40°C) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.