

YaraVita Coprac™ Dual

APPLICATION RATES AND TIMINGS

Avocado: 0.5 l/ha applied to flush of new growth prior to flowering.

Water rate: minimum 500 l/ha.

Bananas: One application of 0.5 l/ha at early canopy development before flowering. Water rate: minimum 30 l/ha.

Beans (Field Growth): 0.5 l/ha at the 4 to 6 leaf stage.

Water rate: minimum 30 l/ha.

Carrot: 0.5 l/ha when crop is 15 cm tall.

Water rate: minimum 50 l/ha.

Cereals: 0.5 l/ha from 2 leaf stage to second node detectable (Zadok's G.S. 12 to 32). Water rate: minimum 50 l/ha.

Coffee: 1 l/ha applied at start of new growth.

Water rate: minimum 400 l/ha.

Eucalyptus: 0.5 to 3 l/ha applied during spring or autumn flush.

Water rate: minimum 20 l/ha.

Garlic: 0.5 l/ha two weeks after transplanting, or in the case of direct sown crops, when the crop is 15 cm tall. Water rate: minimum 75 l/ha.

Groundnuts/Peanuts: 0.5 l/ha at the 4 to 6 leaf stage.

Water rate: minimum 200 l/ha.

Maize, Sorghum: 0.5 l/ha at the 4 to 8 leaf stage.

Water rate: 30 to 200 l/ha.

Nursery Stock/Ornamentals: 0.25 l/ha as soon as there is sufficient leaf area to intercept a spray. Repeat at 10 to 14 day intervals as necessary. Do not spray once first buds have started to open. Spray a maximum of three applications per crop per annum. Water rate: minimum 100 l/ha. N.B. This product may leave a visible residue.

Onion: 0.5 l/ha two weeks after transplanting, or in the case of direct sown crops, when the crop is 15 cm tall. Repeat applications at 10 to 14 day intervals. Water rate: minimum 50 l/ha.

Peas, Pepper (Field Growth): 0.25 l/ha at the 4-6 leaf stage.

Water rate: minimum 200 l/ha.

Potatoes: 0.5 l/ha applied 7 to 14 days after 100% emergence.

Water rate: minimum 200 l/ha.

Rice: 0.5 l/ha applied at start of tillering. Water rate: minimum 200 l/ha

Sugar Cane: 0.5 l/ha when cane is between 30 and 120 cm tall.

Repeat applications may be necessary at 10 to 14 day intervals.

Water rate: minimum 30 l/ha.

Sunflower: 0.25 l/ha applied at 4 to 5 leaf stage.

Water rate: minimum 200 l/ha

Tea: 0.5 l/ha applied during plucking season. Repeat at approximately 1 month intervals. Spray 2 to 6 applications.

Water rate: minimum 200 l/ha.

Tobacco: 0.5 l/ha at 3 to 4 leaf stage. Repeat 10 days later if necessary

Water rate: minimum 30 l/ha.

Tomato (Field Growth): 0.5 l/ha at the 4 to 6 leaf stage.

Water rate: minimum 200 l/ha.

SEED DRESSING

Cereals (barley, wheat) Maize, Rice: 4 litres per tonne.

For further crop recommendations contact your local Yara representative

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 40°C) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.