

YaraVita Tracel™ BZ

APPLICATION RATES AND TIMINGS

Coffee: 2 kg/ha applied early in the growing season. Repeat at the same rate at the pre-flowering stage and also post-harvest.
Water rate: 500 to 1000 l/ha.

Bananas, Citrus, Mango: 2 kg/ha applied to new leaf growth in the spring and again 10 to 14 days later. Repeat the application after final harvest. N.B. Do not apply during flowering.
Water rate: 600 to 1000 l/ha.

Beans, Peas: 2 to 4 kg/ha at the 4 to 6 leaf stage. A second application may be made 10 to 14 days later.
Water rate: 200 to 400 l/ha.

Cabbage, Kale: 2 kg/ha at the 4 to 6 leaf stage, with repeat applications at 10 to 14 day intervals. Water rate: 200 to 500 l/ha.

Capsicum (Field Grown), Tomato (Field Grown): 2 kg/ha when the plants are at the 4 to 6 leaf stage. Repeat once or twice at 10 to 14 day intervals. Water rate: 200 to 500 l/ha.

Cereals (Barley, Rice, Wheat): 2 to 4 kg/ha at the 4 to 6 leaf stage. Water rate: 200 to 400 l/ha.

Cocoa / Cacao: 2 kg/ha applied early in the growing season. Repeat at the same rate at the pre-flowering stage and also post-harvest.
Water rate: 500 to 1000 l/ha.

Forestry (Conifers, Eucalyptus): Two applications of 2 to 4 kg/ha when there is new season leaf production.
Water rate: 200 to 1000 l/ha.

Maize, Sorghum: 2 to 4 kg/ha applied from 4 to 6 leaves.
Water rate: 200 to 400 l/ha.

Onion: 1 to 2 kg/ha as soon as there is sufficient foliage to intercept the spray. A second application may be made at the same rate 10 to 14 days later. Water rate: 200 to 500 l/ha.

Pineapple: 2 kg/ha applied to new leaf growth in the spring and again 10 to 14 days later. Repeat the application after final harvest.
Water rate: 200 to 1000 l/ha.

Potatoes: 2 kg/ha 7 to 14 days after full emergence. Repeat at tuber initiation and at tuber bulking. Water rate: 200 to 500 l/ha.

Sugar Cane: 2 kg/ha when the cane is between 30 and 90 cm tall. Repeat the application 10 to 14 days later.
Water rate: 200 to 500 l/ha.

Sunflower, Oilseed Rape: 2 to 4 kg/ha from 4 to 6 leaves. Consider a second application 10 to 14 days later. N.B. Do not apply during flowering. Water rate: 200 to 400 l/ha.

Tea: 2 kg/ha applied to new leaf growth in the spring and again 10 to 14 days later. Repeat the application after final harvest.
N.B. Do not apply during flowering. Water rate: 600 to 1000 l/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 40°C) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

For further crop recommendations contact your local Yara representative