

YaraVita Bortrac 150

APPLICATION RATES AND TIMINGS

FOLIAR APPLICATION

Apple: 1-2 L/ha at open cluster and again at pink/white bud.

Water rate: 500-1000 L/ha. Also, 2-3 L/ha after harvest but before leaf senescence.

Apricot, Cherry, Nectarine, Peach, Plum: 2 L/ha pre-flowering and 2 applications of 2 L/ha applied after petal fall and repeat at 7 to 14 days later. Water rate: 500-2000 L/ha. Autumn: 2-3 L/ha after harvest but before leaf senescence.

Avocado: 4 applications of 2 to 3 L/ha at 30% and 70% cauliflower pre-flower and post-flower. Water rate: 500-2000 L/ha.

Beans, Lupins, Peas: 2-3 L/ha when plants are 10 to 15 cm high or there is sufficient leaf area to intercept spray. Water rate: 50-200 L/ha.

Blackcurrant: 2 applications of 1 L/ha applied at flower bud stage and repeated 10 to 14 days later (start of flowering). Also, apply 2 L/ha post-harvest, pre-leaf senescence. Water rate: 200-500 L/ha.

Brassicas: 3 L/ha at 4 to 6 leaf stage with repeat applications at the above rate at 10 to 14 day intervals for moderate to severe deficiency. Water rate: 50-500 L/ha.

Carrots: 3 to 5 L/ha soil-applied prior to sowing. 3 L/ha foliar-applied when sufficient leaf area to intercept spray, repeated if necessary 10 to 14 days later. Water rate: 50-200 L/ha.

Cereals: 1 L/ha. Apply from the 4 leaf stage to pseudo stem erection, (Zadoks Growth Stage 14 to 30). Water rate: 200 L/ha.

Citrus: 2 to 3 L/ha at white bud or when flower buds separated. Water rate: 500-1000 L/ha.

Cucurbits (Field Grown): 2 L/ha at 4 leaf stage, repeated if necessary 10 to 14 days later. Water rate: 50-200 L/ha.

Grapevines: 1.5 L/ha at flower truss visible and at flower buds separated, and again at fruit set. Water rate: 500-1000 L/ha. Soil application: 3 to 5 L/ha.

Lettuce (Field Grown): 1 to 2 L/ha 10 to 14 days after transplanting or emergence. Water rate: 500 L/ha.

Olive: 2-3 L/ha applied at bud break and again at the opening of the calyx if necessary. In addition, one or two applications of 2-3 L/ha at a 10 to 14 day interval post-flowering. Also, 2-3 L/ha applied post-harvest. Water rate: 400-1000 L/ha.

Onion: 2-4 L/ha soil-applied prior to sowing. Alternatively, 2 L/ha when there is sufficient leaf area to intercept the spray. A second application may be applied at the same rate 10 to 14 days later. Water rate: 50-200 L/ha.

Pear: 1-2 L/ha at open cluster, pink/white bud and again at petal fall. Water rate: 500-1000 L/ha. Also, 2-3 L/ha after harvest but before leaf senescence.

Potatoes: 2 L/ha 10 to 14 days after 100% emergence. Repeat if necessary 10 to 14 days later. Water rate: 50-200 L/ha.

Strawberry (Field Grown): 1.5 L/ha applied at green bud and again at white bud. Water rate: 200-500 L/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THE LABEL INSTRUCTIONS.

Mix the product thoroughly before use. Pour through the top filter into the tank three-quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is completed. *For more approved recommendations please contact your local Yara representative*

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.*

PRECAUTIONS


Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and faceshield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara NZ Ltd, available on request. Use of the product implies that the purchaser accepts these Conditions of Sale.



WLNZ0202
12.06.18