

YaraVita Mantrac Pro

APPLICATION RATES AND TIMINGS

Apple: 1 to 1.5 l/ha prior to flowering in the case of severe deficiency.

Otherwise commence applications 4 weeks after petal fall and repeat 10 to 14 days later. Also, 1 to 1.5 l/ha after harvest before leaf senescence. Water rate: 500-1000 l/ha. For improved green background colour with the variety Granny Smith only: 6 applications of 0.75 l/ha from December drop to around 4 weeks pre-harvest. Water rate: 200 l/ha.

Apricot, Cherry, Nectarines, Peach, Plum: 1-1.5 l/ha applied from 4 weeks after petal fall and repeated if necessary at 10 to 14 day intervals. Also, 1 l/ha after harvest before leaf senescence.

Water rate: 500-1000 l/ha.

Beans, Peas: 1 to 1.5 l/ha from start of flowering to swelling of pods Repeat application if deficiency is severe. Water rate: 50-200 l/ha.

Brassicas, Canola: 1.25-1.5 l/ha from 4 to 9 true leaves.

Water rate: 50-500 l/ha.

Carrots: 1 to 1.5 l/ha when sufficient leaf area to intercept spray and, if necessary, at 7 to 10 day intervals after this.

Water rate: 50-200 l/ha.

Cereals: 1 to 2 l/ha from 2 leaf stage to first node detectable (Zadok's G.S. 12 to 31). For moderate to severe deficiency repeat applications at 10 to 14 day intervals. Water rate: 30-200 l/ha. For aerial applications 0.5 l/ha applied at the timings described above.

Water rate: 20-50 l/ha.

Clover: 1 l/ha just after every cut. Water rate: 50-200 l/ha.

Citrus: 1.25-1.5 l/ha during spring and autumn flushes.

Water rate: 500-1000 l/ha.

Grapevines: 750 ml/ha at flower truss visible, at flower buds separated and at fruit set. Water rate: 500-1000 l/ha.

Kiwi: 1 l/ha during extension shoot development and from 2 weeks post petal fall (avoid flowering). Water rate: 500-2000 l/ha.

Olive: 1 l/ha. Apply at the start of the season. Repeat applications at 10 day intervals if necessary. Avoid flowering.

Water rate: 1000 l/ha.

Onion: 1 l/ha when sufficient leaf area to intercept spray. Water rate: 50-200 l/ha.

PerSIMMONS: 1-1.5 l/ha applied from 4 weeks after petal fall and repeat if necessary at 10 to 14 day intervals. Also, 1 l/ha after harvest before leaf senescence. Water rate: 500-1000 l/ha.

Potatoes: 1 to 2 l/ha at one week after 100% emergence. For moderate to severe deficiency, repeat applications at 10 to 14 day intervals. Also, 1 l/ha following petiole analysis during tuber bulking. Water rate: 30-200 l/ha.

Strawberry (Field Grown): 1 l/ha at green bud. Repeat if needed at 10 to 14 day intervals. Water rate: 200-500 l/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THE LABEL INSTRUCTIONS.

Mix the product thoroughly before use. Pour through the top filter into the tank three-quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is completed. *For more approved recommendations please contact your local Yara representative*

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.*

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and faceshield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara NZ Ltd, available on request. Use of the product implies that the purchaser accepts these Conditions of Sale.

