

Croplift-K

APPLICATION RATES AND TIMINGS

Apple: 5 kg/ha at 10 to 14 day intervals as required commencing 4 weeks after petal fall. If necessary, also apply post-harvest before leaf fall. Water rate: 500-1000 l/ha.

Apricot, Cherry, Nectarines, Peach, Plum: 5 kg/ha. Commence applications at stone hardening and repeat at 10 to 14 day intervals up to one week before harvest. Water rate: 500-1000 l/ha.

Brassicas: 5 kg/ha. Commence application at 4 to 6 true leaves and repeat as necessary at 7 to 10 day intervals between applications. Water rate: 250-750 l/ha.

Carrots: 5 kg/ha. When sufficient leaf area to intercept spray with 10 to 14 day intervals between applications. Water rate: 200-500 l/ha.

Cereals: 5 kg/ha from mid-tillering to 2nd node detectable (Zadoks G.S. 25 to 32). Repeat at 10 to 14 day intervals between these growth stages as necessary. Water rate: 200 l/ha. For Aerial Application: 5 kg/ha from mid-tillering to 2nd node detectable (Zadoks G.S. 25 to 32). Repeat at 10 to 14 day intervals between these growth stages as necessary. Water rate: 50 l/ha.

Citrus: 5 kg/ha. Commence applications after fruit set and repeat as necessary at 14 to 21 day intervals. Water rate: 1000 l/ha minimum.

Cucurbits (Field Grown): Apply 5 kg/ha at 10 day intervals from setting of first fruit/first fruit development. Water rate: 500 l/ha.

Grapevines: 5 kg/ha from fruit set and repeat at 10 to 14 day intervals as required until pre-harvest. Water rate: 500-1000 l/ha.

Kiwi: 5 kg/ha. Applications from 6 to 8 weeks post petal fall and repeat at 10 to 14 day intervals. Water rate: 500-1000 l/ha.

Olive: 5 kg/ha. Apply as and when required during the growing season. Repeat applications can be made at 10 to 14 day intervals. Avoid applications during flowering. Also, apply as necessary post-harvest. Water rate: 1000 l/ha.

Onions: 5 kg/ha. When sufficient leaf area to intercept spray. Water rate: 200-500 l/ha.

Pears: 5 kg/ha at bud burst. Also, two to three applications at 10 to 14 day intervals, starting at petal fall. A further application of 5 kg/ha may be applied post-harvest, before leaf fall. Water rate: 500-1000 l/ha.

Potatoes: 5 kg/ha. Apply during bulking and repeat as necessary at 7 to 10 day intervals. Water rate: 200-500 l/ha.

Raspberry, Strawberry (Field Grown): 5 kg/ha. Commence applications after flowering and repeat as necessary at 7 to 10 day intervals. Water rate: 500-1000 l/ha.

Turf: 5 kg/ha. As soon as growth commences in the spring. In the case of moderate to severe deficiency, repeat applications at 14 day intervals. Water rate: 500-1000 l/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THE LABEL INSTRUCTIONS.

Slowly wash the product through the top filter into the tank three-quarters filled with water, whilst agitating. Top up with water and continue to agitate until spraying is completed.

For more approved recommendations please contact your local Yara representative

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.*

PRECAUTIONS


Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and faceshield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara NZ Ltd, available on request or at www.yaravita.com. Use of the product implies that the purchaser accepts these Conditions of Sale.



5