

YaraVita Coptrel™ 500

APPLICATION RATES AND TIMINGS

FOLIAR APPLICATION

Carrot: 0.5 l/ha when crop is 15 cm tall. For moderate to severe deficiency repeat applications at 10 to 14 day intervals. Water rate: 200 l/ha.

Cereals: 0.5 l/ha from 2 leaf to second node detectable (Zadok's G.S. 12 to 32). For moderate to severe deficiency repeat applications at 10 to 14 day intervals, within the growth stages indicated. Water rate: 200 l/ha

Conifers: 2 applications of 0.5 l/ha, once there is new season leaf production and again in early Autumn. Water rate: 500 l/ha to 1,000 l/ha.

Grass (Amenity): 0.5 l/ha as soon as growth commences in spring and /or following identification of need by analysis. Repeat sprays at 10 to 14 day intervals as necessary. Water rate: 200 l/ha.

Grass (Grazing): 0.5 l/ha applied 15 to 21 days before grazing or cutting. (Do not apply to grass for sheep). Water rate: 200 l/ha.

Grass (Growth): 0.5 l/ha as soon as growth commences in the Spring. In the case of moderate to severe deficiency, repeat applications at 10 to 14 days intervals. Water rate: 200 l/ha.

Lily, Tulip (Field Crown): 0.5 l/ha. Repeat applications at 7 to 14 day intervals commencing when crop is 10 to 15 cm tall. Water rate: 200 l/ha. Do not apply to crops grown under glass or plastic. Do not apply to the crop in flower.

Maize: 0.5 l/ha at the 4 to 8 leaf stage.

Water rate: 200 l/ha.

Onion: 0.5 l/ha two weeks after transplanting, or in case of direct-sown crops, when the crop is 15cm tall. Repeat applications at 10 to 14 day intervals.

Water rate: 200 l/ha.

Parsnip: 0.5 l/ha when crop is 15 cm tall. For moderate to severe deficiency, repeat applications at 10 to 14 day intervals. (3 applications maximum).

Water rate: 200 l/ha.

Potatoes: 0.5 l/ha applied 7 to 14 days after 100% emergence and following petiole analysis during tuber bulking. Water rate: 200 l/ha.

Sugar Beet: 0.5 l/ha at the 4 to 6 leaf stage. For moderate to severe deficiency, repeat applications should be made at 10-14 day intervals. Water rate: 200 l/ha.

Turf: 0.5 l/ha as soon as growth commences in spring and /or following identification of need by analysis.

Repeat sprays at 10 to 14 day intervals as necessary. Water rate: 200 l/ha.

SOIL APPLICATION

All crops: 5 to 10 l/ha pre-drilling or pre-planting (or before re-growth of perennial crops). Water rate: 50 l/ha minimum.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 40°C) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.