

Magflo 300

APPLICATION RATES AND TIMINGS

Apple, Pears: 4 l/ha after petal fall. In cases of severe deficiency apply also before flowering (around pink bud stage).

Water rate: 500-1,000 l/ha.

Beans, Peas: 2-4 l/ha when crop 15 cm tall. For moderate to severe deficiency repeat applications at 10 to 14 day intervals.

Water rate: 200 l/ha.

Broccoli, Brussels Sprouts, Cabbage, Calabrese, Cauliflower:

2-4 l/ha at 4 to 6 leaf stage, with repeat applications at the above rate to 10 to 14 day intervals for moderate to severe deficiency.

Water rate: 200 l/ha.

Carrot: 2-4 l/ha when crop 15 cm tall. For moderate to severe deficiency repeat applications at 10 to 14 day intervals. Note: On carrots, final application to be made one month before harvest. Water rate: 200 l/ha.

Cereals: 2-4 l/ha at 2 leaf stage to first node detectable (G.S. 12-31). For moderate to severe deficiency, repeat applications at 10-14 day intervals.

2 l/ha at flag leaf ligule just visible to anthesis complete (G.S. 39-69).

Water rate: 200 l/ha.

Cherry, Plum: 4 l/ha at fruit set. If necessary a second application may be made 10 to 14 days later. Water rate: 200 to 500 l/ha.

Grass (Grazing): 5 l/ha applied 7 to 14 days before grazing.

Water rate: 200 l/ha.

Leek, Onion: 2 to 4 l/ha two weeks after transplanting, or in the case of direct sown crops, when the crop is 15 cm tall. Repeat applications at 10 to 14 day intervals. Water rate: 200 l/ha.

Lettuce (field grown): 2-4 l/ha 2 weeks after transplanting. For moderate to severe deficiency repeat applications at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest.

Water rate: 500 l/ha.

Maize: 2-4 l/ha at 4-6 leaf stage. Water rate: 200 l/ha.

Oilseed Rape: For a single application, 2-4 l/ha at onset of stem extension. For moderate deficiency, 2-4 l/ha at 4-6 leaf stage and 2-4 l/ha at onset of stem extension. An extra application can be made 10-14 days later for severe deficiency. Avoid flowering.

Water rate: 200 l/ha.

Potatoes: 2-4 l/ha 1 week after 100% emergence and following petiole analysis during tuber bulking. For moderate to severe deficiency, repeat applications at 10-14 day intervals. Water rate: 200 l/ha.

Raspberry: 4 l/ha at start of flowering. Water rate: 200 to 500 l/ha.

Red Beet, Sugar Beet, Swede, Turnip: 2-4 l/ha at 4 to 6 leaf stage, with repeat applications at the above rate at 10 to 14 day intervals for moderate to severe deficiency. Water rate: 200 l/ha.

Strawberry (Field Grown): 4 l/ha at green bud.

Water rate: 200-500 l/ha.

PROTECTED CROPS

0.2 litres per 100 litres water maximum concentration.

Water rate: 1000 l/ha maximum. Refer to equivalent field grown crop for application timing. N.B. Magflo 300 may leave a white deposit. Do not apply within one month of harvest.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.*

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and faceshield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara, available on request. Use of the product implies that the purchaser accepts these Conditions of Sale.