

Magphos K

APPLICATION RATES AND TIMINGS

Beans, Peas: 1-2 applications of 5 l/ha when crop is approximately 10-15 cm tall. Allow 10-14 days between treatments. Water rate: 200 l/ha.

Broccoli, Brussel Sprouts, Cabbage, Calabrese, Cauliflower:

4 l/ha applied soon after transplanting. Water rate: 200 l/ha.

Carrot: 3 to 5 l/ha when crop is 15 cm tall. Repeat if necessary at 10 to 14 day intervals. Water rate: 200 l/ha.

Cereals: 5 l/ha at tillering. Repeat at 10-14 day intervals if necessary. Also apply from ear emergence until the end of flowering (Zadok's G.S. 51 to 69). Water rate: 200 l/ha.

Daffodil, Lily, Tulip (Field Crown): 5 l/ha applied 7 to 21 days after the flowers have been removed, and repeated again 10 to 14 days later. Water rate: 200 l/ha. Do not apply to crops grown under glass or plastic. Do not apply to the crop in flower.

Grass (Amenity): 5 to 10 l/ha as soon as growth commences in spring and/or following identification of need by analysis. Repeat sprays at 10 to 14 day intervals as necessary. Water rate: 200 l/ha

Leek: 5 l/ha applied from two weeks after transplanting, or in the case of direct sown crops when the crop is 15 cm tall. Repeat applications may be necessary at 10 to 14 day intervals. Water rate: 200 l/ha.

Maize: 5 l/ha at 4 to 8 leaf stage. Repeat at 10 to 14 day intervals if necessary. Water rate: 200 l/ha.

Oilseed Rape: For a single application, 5 l/ha at onset of stem extension. For moderate deficiency, 5 l/ha at 4 to 6 leaf stage and at onset of stem extension. An extra application can be made 10 to 14 days later for severe deficiency. Avoid flowering. Water rate: 200 l/ha

Onion: 5 l/ha. Apply when the foliage is 15 cm tall, with a second application 10 to 14 days later if necessary. Also, 1 to 2 applications of 5 l/ha during bulb filling, with a 10 to 14 day interval between sprays. Water rate: 200 l/ha.

Parsnip: 5 l/ha when crop is 15 cm tall. For moderate to severe deficiency, repeat applications at 10 to 14 day intervals. (3 applications maximum). Water rate: 200 l/ha.

Plum: 2 to 5 applications of 3 to 5 l/ha from petal fall. Repeat applications at 10 to 14 day intervals. Water rate: 500 to 1000 l/ha.

Potatoes: To increase tuber number, 10 l/ha at tuber initiation (when 50% of the tip swellings are twice the diameter of the rest of the stolon). To increase tuber size a minimum of 2 applications of 5 l/ha or 1 to 2 applications of 10 l/ha during tuber bulking (as soon as first formed tubers are 10 mm in diameter). Allow 10-14 days between applications. Water rate: 200 l/ha

Sugar Beet: 5 l/ha at 4-6 leaf stage. Repeat applications may be made at 10-14 day intervals. Water rate: 200 l/ha.

Turf: 5 to 10 l/ha as soon as growth commences in spring and/or following identification of need by analysis. Repeat sprays at 10 to 14 day intervals as necessary. Water rate: 200 l/ha.

PROTECTED CROPS

0.3 litres per 100 litres water maximum concentration.

Water rate: 1000 l/ha maximum. Refer to equivalent field grown crop for application timing.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.*

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and faceshield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara, available on request. Use of the product implies that the purchaser accepts these Conditions of Sale.