

Photrel Pro

APPLICATION RATES AND TIMING

Oilseed Rape: 3 kg/ha* For autumn-sown crops, apply prior to winter dormancy and in spring up to flower buds enclosed (G.S. 3,1). For spring-sown crops, apply at 4 to 9 true leaves followed by a further application 10 days later. Water rate: 200 l/ha.

Brassicas (Broccoli, Cabbage, Calabrese, Cauliflower, Chinese Cabbage, Sprouts), Beans, Peas, Lupins:

3 kg/ha* as early in the growing season as possible, provided there is adequate leaf cover to intercept the spray (e.g. 4 to 6 true leaf stage). Water rate: 200 l/ha.

Carrots: Two applications of 3 kg/ha applied when the crop is 15 cm tall and again two weeks later. Water rate: 200 l/ha.

Cereals: 3 kg/ha applied at any time from the 2 leaf stage to 1st node detectable (Zadoks G.S. 12 to 31). Water rate: 50 to 200 l/ha.

Conifers: 3 kg/ha at start of new season leaf production and again in early autumn. Water rate: 500 to 1,000 l/ha.

Leeks, Onions: 3 kg/ha applied 2 weeks after transplanting and again 10 to 14 days later. Water rate: 200 l/ha.

Lettuce (field grown): 3 kg/ha 10 to 14 days after transplanting or emergence. Repeat after 10 to 14 days if necessary. Water rate: 500 l/ha.

Lupins: 3 kg/ha at the 4 to 6 leaf stage. Repeat once or twice at 10 to 14 day intervals if necessary (for winter varieties apply the repeat sprays at spring re-growth). Water rate: 30 to 200 l/ha.

Potatoes: 3 kg/ha applied one week after 100% emergence, with a second application 10 to 14 days later. Water rate: 200 l/ha.

Soya Beans: 3 kg/ha* applied at the 4 to 6 leaf stage. Water rate: 50 to 200 l/ha.

Sugar Beet: 3 kg/ha* from 4 to 6 true leaves. Water rate: 200 l/ha.

* Repeat applications can be made at 10 to 14 day intervals.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables, outside the control of Yara, can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. *Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.*

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and faceshield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. *Protect from frost and other climatic extremes. (temperatures exceeding 40°C)*

CONDITIONS OF SALE - LIMITED WARRANTY AND LIMITATIONS OF LIABILITY

Yara warrants that the product within the unopened container conforms to the chemical description on this label. The directions on this label must be followed carefully and are based on tests, believed to be reliable. As the occurrence of unusual weather or soil conditions, and the use of this product, are all beyond the control of Yara, no guarantee, expressed or implied, is made as to the effects of such use or the results obtained, and in no event shall the liability of Yara exceed the price of the product used. The purchaser must assume all risks of storage, use or application of this product. Goods are sold subject to the standard Conditions of Sale of Yara, available on request. Use of the product implies that the purchaser accepts these Conditions of Sale.