

YaraVita Caltrac™

APPLICATION RATES AND TIMINGS

Apples*, Pears: Repeat applications of 1-2 quarts/acre at 7 to 10 day intervals applied from petal fall. A single application of up to 2-4 quarts/acre may be made from petal fall to just after "June drop". Minimum water rate: 20 gallons/acre. Note: Late applications may leave a deposit on the crop. For use in

Apricots*, Cherries*, Nectarines, Peaches, Plums: Regular applications of 1-2 quarts/acre at 7 to 14 day intervals from petal fall. Note: Late applications may leave a deposit on the crop. Under conditions conducive to fruit russet (variety, weather conditions, etc) delay applications until 6 weeks after petal fall. Water rate: 15 to 50 gallons/acre.

Citrus: 1-2 quarts/acre beginning from fruit set at 10 to 14 day intervals. Water rate: 25 to 100 gallons/acre. Note: Late applications may leave a deposit on the crop.

Courgettes, Cucumber, Melons, Zucchini, Squash, (Field Grown): Repeat applications of 1-2 quarts/acre at 7 day intervals during fruit development. Note: Late applications may leave a deposit on the crop. Water rate: 20 to 50 gallons/acre.

Cranberry*: 1-3 pints/acre at early bloom. Water rate: 50 gallons/acre. For use in Wisconsin: This product can be applied in Wisconsin where supplemental calcium has been shown to be of benefit to plant storage organs and where soil test calcium is low. Also on crops where supplemental calcium has been shown to improve the resilience to physiological disorders.

Groundnuts: 1-2 quarts/acre at the 4 to 6 leaf stage followed by a second application 10 to 14 days later. Water rate: 3 to 20 gallons/acre.

Nuts (deciduous): 1-2 quarts/acre 7 days after petal fall. Water rate: 50 gallons/acre. Water rate: 50 gallons/acre.

Onions: 1 to 2 quarts/acre applied at the 6 leaf stage. Repeat if necessary 10 to 14 days later. Also, 2 quarts/acre at bulb swelling, repeated once or twice at 10 to 14 day intervals.

Water rate: 3 to 20 gallons/acre.

Peppers (field grown): Repeat applications of 1-2 quarts/acre from the 4-6 leaf stage. Allow 7 days between applications. Note: Late applications may leave a deposit on the crop. Water rate: 5 to 50 gallons/acre.

Potatoes: 2-3 applications of 1-2 quarts/acre commencing at tuber initiation (when 50% of the tip swellings are twice the diameter of the rest of the stolon) and following petiole analysis during tuber bulking with 10 to 14 day intervals between applications. Water rate: 5 to 50 gallons/acre.

Strawberry (Field Grown): 1-2 quarts/acre applied three times from start of new season leaf growth at 10 to 14 day intervals. Note: Late applications may leave a deposit on berries. Also 1-2 quarts/acre applied as required after final harvest for the season at 10 to 14 day intervals. Water rate: 20 to 50 gallons/acre.

Sweet Potatoes: 2-3 applications of 1-2 quarts/acre commencing at tuber initiation with 10 to 14 day intervals between applications. Water rate: 5 to 50 gallons/acre.

Tomatoes (Field Grown): Repeat applications of 1-2 quarts/acre from the 4 to 6 leaf stage. Allow 7 days between applications. Note: Late applications may leave a deposit on the crop. Water rate: 5 to 50 gallons/acre.

Vines: Three applications of 1-2 quarts/acre at cluster stretch, bunch closure (pea sized berries), start of ripening and two weeks later. Water rate: 20 gallons/acre. Late applications may leave a deposit on berries.

*** For use in Wisconsin:** Caltrac can be applied to this crop in Wisconsin where supplemental calcium has been shown to be of benefit to plant storage organs and where soil test calcium is low. Also on crops where supplemental calcium has been shown to improve the resilience to physiological disorders.

Extended recommendations available at
www.yara.us/agriculture/products/yaravita

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and Reproductive Harm –www.P65Warnings.ca.gov