

# YaraVita™ Ferrichel Ninety™

## APPLICATION RATES AND TIMINGS

**Apples, Pears:** 1 to 3 lb/acre as soon as there is sufficient leaf cover to intercept the spray, or as soon as first symptoms appear. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering.

Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre.

**Apricots, Cherries, Nectarines, Peaches, Plums:** 1 to 3 lb/acre as soon as there is sufficient leaf cover to intercept the spray, or as soon as first symptoms appear. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre

**Citrus:** 1 to 3 lb/acre during Spring and Autumn flushes, at 10 to 14 day intervals as necessary. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre.

**Conifers:** 1 to 3 pounds/acre at start of new season leaf production and again in early autumn. Water rate: 50 to 100 gallons/acre.

**Cucurbits (Field Grown):** 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.

**Lettuce (field grown):** 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.

**Nuts (deciduous):** 1 to 3 lbs/acre as soon as there is sufficient leaf cover to intercept the spray, or as soon as first symptoms appear. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre.

**Melon, Water Melons (Field Grown):** 1 to 3 lbs/acre from two weeks after transplanting, or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre

**Onions:** 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.

**Peppers, Tomatoes (Field Grown):** Peppers (field grown): 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.

**Soybeans:** 1 to 3 lbs/acre from 2 weeks after emergence, or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest. Do not apply during flowering. Water rate: 20 gallons/acre.

**Sugar Cane:** 1-3 lb/acre when plants are 12" tall and again 10 to 14 days later. Water rate: 20 gallons/acre.

**Sweet Corn:** 1 to 3 lb/acre from 2 weeks after transplanting or as soon as there is sufficient leaf cover to intercept the spray. Repeat as necessary at 10 to 14 day intervals. Note: Final application to be made at least one month before harvest. Water rate: 20 gallons/acre.

**Vines:** 1 to 3 lb/acre as soon as there is sufficient leaf cover to intercept the spray, or as soon as first symptoms appear. Repeat as necessary at 10 to 14 day intervals. Do not apply during flowering. Note: Final application to be made at least one month before harvest. Water rate: 50 gallons/acre.

*Extended recommendations available at*

<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

## DIRECTIONS FOR USE

### ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

## TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.

## PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

## STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

## CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



**WARNING:** Cancer and Reproductive Harm –[www.P65Warnings.ca.gov](http://www.P65Warnings.ca.gov)