

YaraVita™ Magphos™

APPLICATION RATES AND TIMINGS

Apples, Pears: 4 to 6 applications of 1-2 quarts/acre from end of flowering. Repeat applications at 10 to 14 day intervals through color break. Water rate: 50 to 100 gallons/acre.

Apricots, Cherries, Nectarines, Peaches, Plums: 2 to 5 applications of 1-2 quarts/acre from petal fall. Repeat applications at 7 to 14 day intervals. Water rate: 50 to 100 gallons/acre.

Blueberries: 1-2 quarts/acre 7-10 days after petal fall has finished. Repeat in 10-14 day intervals through berry softening and color break. Water rate: 100 gallons/acre.

Cereals: 1-2 quarts/acre at tillering. Repeat at 10 to 14 day intervals if necessary. Also apply from ear emergence until the end of flowering (Zadok's G.S. 51 to 69). Water rate: 5 to 20 gallons/acre.

Cole Crops, Brussel Sprouts, Broccoli, Cabbage, Calabrese, Cauliflower: 1-2 quarts/acre applied soon after transplanting. Repeat 2-3 times in 10-14 day intervals during stem extension and head development. Water rate: 7.5 to 20 gallons/acre.

Corn, Sorghum: 1-2 quarts/acre at 4 to 8 leaf stage. Repeat at 10 to 14 day intervals if necessary. Water rate: 3 to 20 gallons/acre.

Cotton: 1-2 quarts/acre at the start of flowering. Repeat in 14 day intervals as required. Water rate: 5 to 20 gallons/acre.

Cucurbits, Squash, Zucchini (Field Grown): 1-2 quarts/acre at the 4 to 6 leaf stage. Repeat at 10 to 14 day intervals. Water rate: 20 to 50 gallons/acre.

Lettuce (field grown): 1-2 quarts/acre. Apply 10 to 14 days after transplanting or emergence. One or two repeat applications may be made at 10 to 14 day intervals. Water rate: 50 gallons/acre.

Melons (Field Grown): 1-2 quarts/acre at start of flowering. Repeat at 10 to 14 day intervals if necessary. Water rate: 50 gallons/acre.

Nuts (deciduous): Two to three applications of 1-2 quarts per acre beginning at petal fall. The repeat applications should be made at 7 to 14 day intervals through hull split. Water rate: 50 to 100 gallons per acre.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

Onions: 1-2 quarts/acre. Apply when the foliage is 6" tall, with repeat applications 10 to 14 day intervals if necessary. Water rate: 3 to 20 gallons/acre.

Peas: 1 to 2 applications of 1-2 quarts/acre when the crop is 4 to 6" tall. Allow 10 to 14 days between applications. Water rate: 3 to 20 gallons/acre.

Peppers (Field Grown): 1 pint in 45 gallons before transplanting. Repeat applications in 10-14 day intervals at 1-2 Quarts per acre. Water rate: 100 gallons/acre maximum.

Potatoes: To increase tuber size: A minimum of 2 applications of 1-2 quarts/acre during tuber bulking (as soon as first formed tuber are 1/2" in diameter) and following petiole analysis during tuber bulking. Allow 10 to 14 days between applications. Water rate: 5 to 20 gallons/acre.

Strawberries (Field Grown): Non everbearing varieties: 3 applications of 1-2 quarts/acre from start of flowering. Repeat applications 7 to 10 day intervals. Everbearing varieties: 6 applications of 1-2 quarts/acre. Do not apply successive applications at intervals of less than 10 to 14 days. Water rate: 30 to 50 gallons/acre.

Sweet Potato: 1-2 quarts/acre one week after 100% emergence or transplanting. Repeat applications during tuber bulking at 10 to 14 day intervals. Also, apply at the same rate following recommendation from analysis. Water rate: 20 gallons/acre.

Tomatoes (Field Grown): 1-2 quarts/acre. Apply at the 4 to 6 leaf stage with repeat applications at 10 to 14 day intervals if necessary. Water rate: 3 to 50 gallons/acre.

Vines: 2 to 5 applications of 1-2 quarts/acre at 10-14 day intervals beginning at early shoot growth, flower buds separated and stretching, after berry set, berry size and early veraison before harvest. Water rate: minimum 20 gallons/acre.

Extended recommendations available at

<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>



WARNING: Cancer and Reproductive Harm –www.P65Warnings.ca.gov