

YaraVita™ Coptrac™

APPLICATION RATES AND TIMINGS

Alfalfa: 0.5 pint/acre applied at start of tillering.

Water rate: 20 gallons/acre.

Carrots: ½ pint/acre when crop is 6" tall. For moderate to severe deficiency, repeat the application once or twice at 10 to 14 day intervals. Water rate: 20 gallons/acre.

Cereals: ½ pint/acre from 2 leaf stage to second node detectable (Zadok's G.S. 12 to 32). For fall-planted crops, an application prior to winter dormancy is strongly recommended. For moderate to severe deficiency, repeat the application once or twice at 10-14 day intervals. Water rate: 20 gallons/acre.

Conifers: 2 applications of ½ pint/acre once there is new season leaf production and again in early autumn.

Water rate: 50 to 100 gallons/acre.

Corn: ½ pint/acre at 4 to 6 leaf stage.

Water rate: 20 gallons/acre.

Garlic: ½ pint/acre two weeks after transplanting, or in the case of direct sown crops, when the crop is 6 inches tall. Repeat applications at 10 to 14 day intervals.

Water rate: 7.5 to 20 gallons/acre.

Nursery/Ornamentals: One pint in 100 gallons water (0.125% v/v) as soon as there is sufficient leaf area to intercept a spray.

Repeat at 10 to 14 day intervals as necessary. Avoid applications during flowering. Spray a maximum of three applications per crop per annum. Note: Do not apply within one month of picking/marketing.

Maximum water rate: 20 gallons/acre.

Onions: ½ pint/acre two weeks after transplanting, or in the case of direct sown crops, when the crop is 6" tall. For moderate to severe deficiency, repeat the application once or twice at 10 to 14 day intervals. Water rate: 20 gallons/acre.

Potatoes: 0.5 pint/acre applied 7 to 14 days after 100% emergence. Water rate: 20 gallons/acre.

Rice: ½ pint/acre applied at start of tillering.

Water rate: 20 gallons/acre.

Sorghum: ½ pint/acre at the 4 to 8 leaf stage.

Water rate: 3 to 20 gallons/acre.

Sweet Corn: ½ pint/acre at 4-6 leaf stage.

Water rate: 20 gallons/acre.

SOIL APPLICATION

All Crops: 2 quarts/acre to 1 gallon/acre, applied pre-planting or pre-emergence. Water rate: 5 gallons/acre minimum

Extended recommendations available at
<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and reproductive Harm - www.P65Warnings.ca.gov