

YaraVita™ Mancozin™

APPLICATION RATES AND TIMINGS

Alfalfa: 1½ pints/acre at dormancy break. Repeat after every cut when re-growth has occurred, as necessary.

Water rate: 3 to 20 gallons/acre.

Beans, Dry Bean, Peas: 1 pint/acre at the 4 to 6 leaf stage (for deficiency/yield). Water rate: 20 gallons/acre.

Cole Crops, Brussel Sprouts, Cabbage: 1 pint/acre at the 4 to 6 leaf stage. Repeat 10 to 14 days later if necessary.

Water rate: 20 gallons/acre.

Canola: 1 pint/acre at the 4 to 6 leaf stage. Repeat 10 to 14 days later if necessary. Water rate: 20 gallons/acre.

Cereals: 1 pint/acre at two leaf stage to first node detectable (Zadok's G.S. 12-31). For severe deficiency, increase to 1 quart/acre or repeat at the above rate. For autumn sown crops, an application prior to winter dormancy is strongly recommended. Water rate: 20 gallons/acre.

Garlic: 1 pint/acre two weeks after transplanting, or in the case of direct sown crops, when the crop is 6 inches tall. Repeat applications at 10 to 14 day intervals.

Water rate: 20 gallons/acre.

Citrus: 1 to 2 pints/acre. During spring and autumn flushes.

Water rate: 50 to 200 gallons/acre.

Corn: 1 pint/acre at the 4 to 8 leaf stage. For moderate to severe deficiency repeat applications at 10 to 14 day intervals.

Water rate: 3 to 20 gallons/acre.

Grapevines: 1 to 1½ pints/acre applied at 10 to 14 day intervals pre-and post- flowering. Avoid applications during flowering.

Water rate: 20 to 100 gallons/acre.

Lettuce (Field Grown): 1 pint/acre at the 4 to 6 leaf stage.

Water rate: 20 gallons/acre.

Melon (Field Grown): 1 pint/acre applied at the 4 to 6 leaf stage. In cases of moderate or severe deficiency repeat the application after a 10 to 14 day interval.

Water rate: 50 gallons/acre.

Nuts (Deciduous): 1 pt./acre at bud break and 10 to 14 days later. N.B. Do not apply during flowering. Also, 1pt/acre after harvest but before leaf senescence.

Water rate: 20 to 100 gallons/acre.

Onion: 1½ quarts/acre when there is sufficient leaf area to obtain coverage (4 to 6 leaf). Repeat during bulking if necessary. Water rate: 3 to 20 gallons/acre.

Rice: ½ to 1 pint/acre at the start of tillering.

Water rate: 3 to 20 gallons/acre.

Sorghum: 1 pint/acre at the 4 to 8 leaf stage. Repeat if necessary at 10 to 14 day intervals.

Water rate: 3 to 20 gallons/acre.

Soya Bean: 1 pint/acre when crop is 2 to 6 inches tall. Repeat if necessary 10 to 14 days later. Water rate: 3 to 20 gallons/acre.

Tomato (Field Grown): 1 to 2 pints/acre at the 4 to 6 leaf stage. Repeat when required at 10 to 14 day intervals.

Water rate: 3 to 150 gallons/acre.

FERTILIZER IMPREGNATION

Mancozin may be impregnated onto dry fertilizer granules except those which contain or are based on ammonium nitrate. Consult your YARA technical representative for details.

Extended recommendations available at

<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (Temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and reproductive Harm - www.P65Warnings.ca.gov