

YaraVita™ Agripotash™

WARNING

This fertilizer is to be used only on crops which respond to molybdenum (Mo). Crops high in molybdenum (Mo) are toxic to grazing animals (ruminants).

APPLICATION RATES AND TIMINGS

Apples, Pears: 2 quarts/acre at bud burst and also post-harvest before leaf fall. Water rate: 50 gallons/acre.

Apricots, Cherries, Nectarines, Peaches: 3 applications of 1½ to 2 quarts/acre from stone hardening, with repeat applications at 10 to 14 day intervals. Water rate: 50 to 100 gallons/acre.

Canola: 2 quarts/acre at the 4 to 6 leaf stage. Repeat as required for moderate to severe deficiency at 7 to 14 day intervals. Water rate: 20 gallons/acre.

Cereals: 2 quarts/acre at first node detectable (Zadok's G.S. 31). Water rate: 20 gallons/acre.

Cole Crops, Broccoli, Brussel Sprouts, Cabbage, Calabrese, Cauliflower: 2 quarts/acre at the 4 to 6 leaf stage. Repeat as required for moderate to severe deficiency at 7 to 14 day intervals. Water rate: 20 gallons/acre.

Cotton: 2 quarts/acre at squaring and at flowering. Water rate: 20 gallons/acre.

Courgette, Cucumber, Squash, Water Melons, Zucchini (Field Grown): Apply 1 to 2 quarts/acre at 10 day intervals from setting of first fruit/first fruit development. Water rate: 50 gallons/acre.

Corn: 2 quarts/acre at the 4 to 8 leaf stage. Water rate: 20 gallons/acre.

Melons (Field Grown): 2 quarts/acre applied from first fruit development. Repeat at 10 day intervals.

Water rate: 50 gallons/acre.

Nuts (Deciduous): 2 quarts/acre at bud break in spring. Water rate: 50 gallons/acre.

Onions: 2 quarts/acre when sufficient leaf area to intercept spray. Water rate: 20 to 50 gallons/acre.

Potatoes: 2 applications of 2 quarts/acre during tuber bulking (as soon as first formed tubers are half an inch in diameter). Allow 10 to 14 days between applications.

Water rate: 20 gallons/acre.

Raspberry, Strawberries (Field Grown): 2 quarts/acre at green bud. Water rate: 50 gallons/acre.

Soybeans: 1 quart/acre when the crop is 4 to 6 inches tall. Water rate: 20 gallons/acre.

Strawberries (Field Grown): 2 quarts/acre at green bud. Water rate: 50 gallons/acre.

Sugar Cane: 2 quarts/acre when cane is between 12 and 50 inches tall. Repeat applications may be necessary at 10 to 14 day intervals. Water rate: 3 to 20 gallons/acre.

Sweet Potatoes: 2 quarts/acre one week after 100% emergence or transplanting. Also, apply at the same rate following recommendation from analysis. Water rate: 20 gallons/acre.

Tobacco: Three applications of 1½ to 2 quarts/acre two to three weeks after transplanting (3 to 4 leaf stage) with 10 days between applications. Water rate: 40 gallons/acre.

Tomatoes (Field Grown): 1½ quarts/acre applied at first truss, fruit set and repeated at 10 day intervals. Water rate: 50 gallons/acre.

Vines (Wine Grapes Only): 1½ to 2 quarts/acre at fruit set, pea-sized berries and first color softening/one month before harvest. Water rate: 50 gallons/acre.

Extended recommendations available at

www.yara.us/agriculture/products/yaravita

WARNING: This product may contain a chemical (or chemicals) known to the State of California to cause cancer and birth defects or other reproductive harm.

DIRECTIONS FOR USE ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.