

YaraVita™ Seniphos™

APPLICATION RATES AND TIMINGS

Almond, Nuts (Deciduous): 2 to 5 applications of ½ - 1 gallon/acre at 7 to 14 day intervals commencing at petal fall. Water rate: 50 to 100 gallons/acre.

Apples: 3 to 8 applications of 1-4 quarts/acre at 10 to 14 day intervals commencing at petal fall. One or two applications (7 days apart) should be applied alone once the fruit has started to change colour, normally 2 to 3 weeks before harvest. Water rate: 50 to 100 gallons/acre.

Apricots, Cherries, Nectarines: 2 to 5 applications of 1-4 quarts/acre at 7 to 14 day intervals commencing at petal fall. Water rate: 50 to 100 gallons/acre.

Blueberries: 2-4 quarts/acre applied 7-10 days after petal fall. Repeat during berry sizing. Water rate: 100 gallons/acre.

Broccoli, Brussel Sprouts, Cabbage, Calabrese, Cauliflower, Cole: 2 to 3 applications of 1-4 quarts/acre from stem extension/head development at 7 to 14 day intervals. Water rate: 7.5 to 20 gallons/acre.

Citrus: 4 quarts/acre at fruit set with 1 to 2 further applications at 10 to 14 day intervals. Water rate: 50 to 100 gallons/acre.

Corn: 2 quarts/acre at 4 to 8 leaf stage. Water rate: 20 gallons/acre.

Cotton: 2 quarts/acre from 10% flowering at 14 day intervals. Water rate: 30 gallons/acre.

Cucurbits (Field Grown): 3 to 5 applications of 2 quarts/acre at 7 day intervals commencing at fruit set. Water rate: 30 gallons/acre.

Garlic, Onions: 1 to 2 applications of 2 quarts/acre during bulb filling, with a 10 to 14 day interval between sprays. Water rate: 20 gallons/acre

Lettuce (Field Grown): 2 to 3 applications of 2 quarts/acre commencing 10 to 14 days after transplanting or emergence with 7 to 10 day intervals between applications. Water rate: 20 to 50 gallons/acre

Melons (Field Grown): 1½ to 2 quarts/acre at start of flowering. Repeat at 10 to 14 day intervals if necessary. Water rate: 50 gallons/acre.

Peaches, Plums: 2 to 5 applications of 2-4 quarts/acre at 7 to 14 day intervals commencing at petal fall. Water rate: 50 to 100 gallons/acre. For red coloration: Seniphos at 4 quarts/acre in 100 gallons of water has been shown to enhance red coloration. Seniphos should be applied Alone and NOT mixed with agrichemicals. One or two applications (7 days apart) should be made once the fruit has started to change colour, normally 2 to 3 weeks before harvest.

Pears: 3 to 8 applications of 1- 4 quarts/acre at 10 to 14 day intervals commencing at petal fall. Water rate: 50 gallons/acre.

Potatoes: A minimum of 2 applications of 2 to 4 quarts/acre during tuber bulking (as soon as first formed tubers are ½" in diameter) and following petiole analysis during tuber bulking. Allow 10 to 14 days between applications. Water rate: 20 gallons/acre.

Raspberry: 3 applications of 1- 4 quarts/acre. Start of flowering, end of flowering and fruit development. Water rate: 50 gallons/acre

Strawberries (Field Grown): Non-everbearing varieties: 3 applications of 1- 4 quarts/acre from start of flowering. Repeat applications at 7 to 10 day intervals. Everbearing varieties: Divide a total rate of 12 quarts/acre into 6 applications of 2 quarts/acre. Do not apply successive applications at intervals of less than 10 to 14 days. Water rate: 20 to 50 gallons/acre. Strawberries (field grown in Florida): Regular applications of 1 to 2 quarts/acre from start of flowering. Repeat applications at 7 to 10 intervals. Water rate: 20 to 50 gallons/acre.

Tomatoes (Field Grown): 2 to 4 applications of 2 quarts/acre commencing from flowering on second truss. Repeat at 10 to 14 day intervals. Water rate: 50 gallons/acre.

Vines: 1-4 quarts/acre at fruit set and repeated at 10 to 14 day intervals till color break. Water rate: 50 gallons/acre.

Extended recommendations available at
<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

DIRECTIONS FOR USE ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (Temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and Reproductive Harm –www.P65Warnings.ca.gov