

YaraVita™ Stopit™

APPLICATION RATES AND TIMINGS

Apple*: Regular applications of 2-4 quarts/acre at 7 to 14 day intervals from petal fall to one week before harvest.

Water rate: 15 to 50 gallons/acre.

Apricot, Cherry, Nectarines, Peach, Plum: 2 applications of 2-4 quarts/acre applied 4 weeks and 2 weeks before harvest.

Water rate: 15 to 50 gallons/acre.

Cole Crops, Broccoli, Brussel Sprouts, Cabbage, Calabrese,

Cauliflower: 2 to 3 applications of 2 quarts/acre from stem extension/head development to harvest. Allow 10 to 14 day intervals between applications. Water rate: 7.5 to 20 gallons/acre.

Carrots: 2 to 3 applications of 2 quarts/acre from when the crop is 4 to 6 inches tall. Allow 10 to 14 day intervals between applications.

Water rate: 20 gallons/acre.

Citrus: 3 applications of 4 quarts/acre applied during the period from fruit expansion to harvest. Allow 10 to 14 day intervals between applications. Water rate: 50 to 100 gallons/acre.

Cucurbits (Field Grown): 2 to 3 applications of 2 quarts/acre commencing at fruit set with 7 day intervals between applications.

Water rate: 20 gallons/acre.

Garlic: ½ to 1 gallon/acre at bulb swelling. Repeat applications at 7 to 14 day intervals as necessary. Water rate: 5 to 20 gallons/acre.

Lettuce (Field Grown): 2 to 3 applications of 1-2 quarts/acre commencing 10 to 14 days after transplanting or emergence with 10 to 14 day intervals between applications.

Water rate: 5 to 50 gallons/acre.

Ornamentals: 1-3 pints/acre as soon as there is sufficient leaf area to intercept a spray. Repeat at 10 to 14 day intervals as necessary. Avoid applications during flowering. Spray a maximum of three applications per crop per annum. Water rate: 10 to 20 gallons/acre.

Onion: 2-4 quarts/acre at bulb swelling. Repeat applications at 7 to 14 day intervals as necessary. Water rate: 5 to 20 gallons/acre

Pears: 2 to 3 applications of 2-4 quarts/acre at 10 to 14 day intervals timed so that the final application is made one week before harvest.

Water rate: 15 to 50 gallons/acre.

Pepper (Field Grown): One to four applications of 1-2 quarts/acre from flowering on second truss onwards.

Water rate: 5 to 50 gallons/acre.

Potatoes*: 2 to 3 applications of 2-4 quarts/acre commencing at tuber initiation (when 50% of the tip swellings are twice the diameter of the rest of the stolon) and following petiole analysis during tuber bulking with 10 to 14 day intervals between applications.

Water rate: 5 to 20 gallons/acre.

Small grains: 2 quarts/acre. Apply from 4 leaves to second node detectable. (Zadok's G, S, 14 to 32). Water rate: 20 gallons/acre.

Strawberry* (Field Grown): Non-everbearing varieties: 3 applications of 2-4 quarts/acre from start of flowering. Repeat applications at 7 to 10 intervals. Water rate: 20 to 50 gallons/acre. Everbearing varieties: Divide a total rate of 12 quarts/acre into 6 applications of 2 quarts/acre. Do not apply successive applications at intervals of less than 10 to 14 days. Water rate: 20 to 50 gallons/acre.

Tomato (Field Grown): Applications of 1-2 quarts/acre commencing from flowering on second truss. Allow 7 days between applications. Water rate: 5 to 50 gallons/acre.

Vines: 2-4 quarts/acre repeated applications at 10 to 14 day intervals from start of colouring. Water rate: 50 to 100 gallons/acre.

***Wisconsin:** Stopit can be applied to this crop in Wisconsin where supplemental calcium has been shown to be of benefit to plant storage organs and where soil test calcium is low. Also on crops where supplemental calcium has been shown to improve the resistance to disease.

Extended recommendations available at

<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and Reproductive Harm –www.P65Warnings.ca.gov