

YaraVita™ Bud Builder WP™

WARNING

This fertilizer contains boron which may be injurious to certain crops. Contact your county agent or field consultant for specific information.

APPLICATION RATES AND TIMINGS

Apples, Pears: In the Spring apply 5 to 10 lbs/acre applied at bud burst. Reapply after fruit set during sizing in 10 to 14 day intervals. 5-10 lbs/acre applied post-harvest before leaf senescence. Water rate: minimum 25 gallons/acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

Apricots, Nectarines Peaches, Cherries, Prunes: 5-10 lbs/acre. Apply from bud burst to start of flowering. Apply petal fall thru early fruit sizing in 10-14 day intervals thru green fruit sizing & leaf canopy development period. Apply post-harvest in 21-28 day interval till leaf senescence. Water rate: minimum 25 gallons/acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

Blackberry, Blueberry, Boysenberry: Apply 3-5 lbs/acre: Apply from bud swell thru pink bud. Apply post bloom-petal fall thru green berry sizing and leaf development period in 14 day intervals.

Reapply post-harvest during fruit bud formation, in 21-28 day interval until the onset of leaf senescence. Water rate: minimum 20 gallons/acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

Brassicas: 5-10 lbs per acre at the 4 to 6 leaf stage, with repeat applications at 10 to 14 day intervals. Water rate: minimum 20 gallons per acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

Citrus, Oranges: 5-10 lbs/acre applied post-harvest. Also, 5-10 lbs/acre applied from bud burst to start of flowering Reapply from petal fall thru fruit sizing in 14-21 day interval. Water rate: minimum 25 gallons per acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

Conifers: 2 applications of 5-10 pounds/acre once there is new season leaf production and again in early autumn. Water rate: 50 to 100 gallons/acre.

Grapevines: Apply 1 to 3 applications of 5-10 lbs/acre applied bud break thru pre-bloom at a 10-14 day interval. Also, during post-bloom thru berry sizing 1-2 applications in 7-14 day intervals. Also apply post-harvest before leaf senescence. Water rate: minimum 25 gallons/acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

Nuts (Deciduous), Almonds, Hazelnuts, Pistachios, Pecans, Walnuts: Apply 5 to 10 lbs/ac applied from bud burst to start of flowering. After nut set, repeat in 14-28 day intervals. Apply 5-10 lbs/ac post-harvest before leaf senescence. Water rate: 25 gal/ac minimum.

Potatoes: 5 lbs/acre one week after plant emergence. Repeat at 10 to 14 day intervals. Water rate: 20 gallons per acre.

Raspberry, Strawberry: 5 lbs/acre applied from green tip to pink bud to just prior to start of flowering in 10-14 day intervals. Apply 1 to 3 times from petal fall thru berry sizing. Also, 5 lbs/acre post-harvest in 21 to 28 day interval up to leaf senescence. Water rate: minimum 20 gallons/acre. Foliar application of Bud Builder may leave a white residue-deposit on the leaf or crop when applied close to harvest.

Extended recommendations available at
<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

DIRECTIONS FOR USE ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and reproductive Harm - www.P65Warnings.ca.gov