

YaraVita Biotrac™

WARNING

This product contains boron which may be injurious to certain crops. Contact your county agent or field consultant for specific information.

APPLICATION RATES AND TIMINGS

Corn: 1 QTS/Acre at the 4 to 8 leaf stage. Water rate: 20 gallons/acre minimum.

Fertigation: 2.5 to 5 QT/acre applied via a fertigation system. Apply alone. Mix and use immediately. Flush system thoroughly immediately after application with water. DO NOT use in "closed" nor in "recirculating" systems, nor apply to plants/crops grown in pots/containers. Repeat at the same rate with the same number of inputs and timing advice as for foliar recommendations for each named crop.

Grapes (table/ wine): 3 pints/acre applied pre-flowering, at flowering and also at fruit set. Water rate: 20 Gal/acre minimum. Alternatively: 3 pints/acre at flower truss visible, at flower buds separated / start of flowering and at fruit set. Water rate: 20 Gal/acre minimum.

Lettuce: Field Grown: Three applications of 3 pints/acre at 10 to 14 day intervals commencing at the 4 to 6 leaf stage. Water rate: 20 GAL/acre minimum.

Melon: 3 pints/acre at 10 to 14 day intervals commencing at the 4 to 6 leaf stage. Water rate: 20 GAL/acre minimum. Continuous Cropping Field Grown Varieties: 3 pints/acre at the 4 to 6 leaf stage. Applications may be repeated at 10 to 14 day intervals, as necessary. Water rate: 200 GAL/acre.

Nuts (deciduous): Apply 3 pints/acre when the first leaves appear, at the start of flowering and at the end of flowering / fruit swelling. Water rate: 20 Gal/acre minimum.

Potatoes: 1 QT/Acre applied 7 to 14 days after 100 percent emergence, 20 days after tuber initiation and, following petiole analysis, during tuber bulking. Water rate: 20 gallons, minimum.

Soya bean: Three applications of 1.25 QTS/Acre at 10 to 14 day intervals commencing at the 4 to 6 leaf stage. Water rate: 20 gallons/acre.

Tomato (Field Grown): Three applications of 1 QT/Acre at 10 to 14 day intervals commencing at the 4 to 6 leaf stage. Water rate: 20 gallons/acre minimum. Continuous Cropping Field Grown Varieties: 1 QT/Acre at the 4 to 6 leaf stage. Applications may be repeated at 10 to 14 day intervals, as necessary. Water rate: 20 gallons/acre.

Extended recommendations available at www.yara.us/agriculture/products/yaravita

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and Reproductive Harm –www.P65Warnings.ca.gov