

# YaraVita Gramitrel™

## APPLICATION RATES AND TIMINGS

**Almond, Pistachios, Walnuts:** 1-2 pints/acre beginning at leaf canopy development 2 weeks after nut set, repeat 14-21 day later if necessary. Additionally, 2-4 pints/acre after harvest before leaf senescence. Water rate: minimum 50 gallons/acre.

**Apricot, Nectarines, Peach, Plum:** 2-4 pints/acre after harvest before leaf senescence. Water rate: minimum 20 gallons/acre.

**Blackberry, Blueberry, Boysen Berry:** 1-2 pints/acre as soon as there is sufficient leaf cover to intercept the spray. Repeat two weeks later if required. Water rate: minimum 20 gallons/acre. Not to apply during flowering or fruit development.

**Cereals:** 1.2 to 4 pints/acre from the 2 leaf stage to second node detectable (Zadok's G.S. 12 to 32). Repeat applications at 10 to 14 day intervals, within this timing, if necessary. Water rate: 5 to 20 gallons/acre.

**Citrus, Grapefruit, Lemon, Oranges:** 3 to 5 pints/acre applied during spring flush and also during summer / autumn flush. Water rate: 50 to 100 Gallons/acre.

**Corn:** 1- 2 pints/acre at 4 to 6 leaf stage. Repeat in 14-21 day interval. Water rate: minimum 20 gallons/acre.

**Grapes (table), Grapevines:** 1 to 2 pints/acre during leaf canopy development pre bloom. Reapply during early fruit sizing at a 10 to 14 day interval and again post-harvest before leaf senescence. Water rate: minimum 20 gallons/acre.

**Grapes (wine):** 1 to 2 pints/acre during leaf canopy development pre bloom. Reapply during early fruit sizing at a 10 to 14 day interval and again post-harvest before leaf senescence. Water rate: minimum 20 gallons/acre.

**Hazelnuts, Pecan:** 1-2 pints/acre beginning at leaf canopy development 2 weeks after nut set, repeat 14-21 day later if necessary. Additionally, 2-4 pints/acre after harvest before leaf senescence. Water rate: minimum 50 gallons/acre.

**Raspberry, Strawberry:** 1-2 pints/acre as soon as there is sufficient leaf cover to intercept the spray. Repeat two weeks later if required. Water rate: minimum 20 gallons/acre. Not to apply during flowering or fruit development.

*Extended recommendations available at*  
<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

## DIRECTIONS FOR USE ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

## TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.

## PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

## STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

## CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



**WARNING:** Cancer and Reproductive Harm –[www.P65Warnings.ca.gov](http://www.P65Warnings.ca.gov)