

YaraVita Safe N™

APPLICATION RATES AND TIMINGS

APPLY ONLY AS DIRECTED.

Almond, Walnuts: 2 to 4 quarts/acre before flowering. Repeat 30 days after flowering, if necessary. Also, after harvest before leaf fall. Water rate: 20 gallons/acre minimum.

Apple: 2 to 4 quarts / acre from bud burst to the start of flowering and/or from petal fall with a repeat application 10 to 14 days later if necessary and/or after harvest before leaf senescence. Water rate: 50 gallons/acre. Alternatively: Apply 4 to 5 quarts/acre prior to fruit set or post harvest. Safe N 25-0-0 can be applied either undiluted or diluted (with water) for aerial or ground application. Use sufficient carrier to adequately cover crop foliage.

Cherry, Peach: 2 to 4 quarts / acre from petal fall with a repeat application 10 to 14 days later if necessary and/or after harvest before leaf senescence. Water rate: 50 gallons/acre. Alternatively; Apply 5 to 9 quarts/acre at early fruit set and repeat at early fruit color. Safe N 25-0-0 can be applied either undiluted or diluted (with water) for aerial or ground application. Use sufficient carrier to adequately cover crop foliage.

Grapes (Table, Wine), Grapevines: Two applications of 2 to 4 quarts/acre at a 10 to 14 day interval from leaves unfurled up to flower buds separated and/or two applications of 4 to 7 quarts/acre at a 10 to 14 day interval from start of veraison. Also, 4 to 6 quarts/acre applied after harvest before leaf senescence. Water rate: 15 gallons/acre minimum.

Lettuce (Field Growth): 4 to 5 quarts/acre. Apply 10 to 14 days after transplanting or thinning. Repeat 14 days later and again 14 days prior to harvest. Water rate: 50 Gallons/acre minimum. Alternatively: 4 to 5 quarts/acre. Apply after thinning, then at

early head formation and repeat at 10 - 14 day intervals. Safe N 25-0-0 can be applied either undiluted or diluted (with water) for aerial or ground application. Use sufficient carrier to adequately cover crop foliage.

Potatoes: 1 to 2 gallons/acre before tuber set and at early bloom. Also, apply 1 to 2 gallons/acre, if required, following petiole analysis. Water rate: 10 to 20 gallons/acre.

Tomato (Field Growth): 1 to 1.5 gallons/acre. Apply at fruit set. Repeat at 10 to 14 day intervals as required. Water rate: 50 gallons/acre minimum. Alternatively: PROCESS: (Single Application) - 2.5 gallons/acre. Apply 10 - 14 days after full bloom. (Multiple Applications) - 1 to 1.5 gallons/acre at full bloom and repeat at 10 - 14 day intervals until harvest is near completion. Safe N 25-0-0 can be applied either undiluted or diluted (with water) for aerial or ground application. Use sufficient carrier to adequately cover crop foliage.

Nursery stock/ornamentals: (Outdoor) 0.5 to 2 gallons Per 1000 gallon of water. Repeat at 10 - 14 day intervals as necessary. Avoid applications during flowering. Safe N 25-0-0 has been applied undiluted to some types of ornamentals and nursery stock. Since these plants vary widely in growth habit and leaf texture, some types are more susceptible to leaf or tip burn than others. It is recommended that the user try only a few plants at undiluted or concentrated rates and observe for a week before general applications are made.

Turf Grass: 3 to 9 fl oz per 1000 sq ft in a water volume of 0.5 to 1.5 galls/1000 sq ft every 7 - 14 days during the growing season based on nutritional requirements.

Extended recommendations available at
<https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and Reproductive Harm –www.P65Warnings.ca.gov